This article was published in an Elsevier journal. The attached copy is furnished to the author for non-commercial research and education use, including for instruction at the author's institution, sharing with colleagues and providing to institution administration.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>

Research paper

Carbonate eolianites, quartz sands, and Quaternary sea-level cycles, Western Australia: A chronostratigraphic approach

Paul J. Hearty^{a,*}, Michael J. O'Leary^b

^a*School of Earth and Environmental Sciences, University of Wollongong, Wollongong, NSW 2522, Australia*

^b*Department of Environmental & Geographical Sciences, Manchester Metropolitan University, Manchester M1 5GD, UK*

Received 16 June 2007; received in revised form 26 September 2007; accepted 3 October 2007

Available online 10 October 2007

Abstract

Marine and eolian carbonate deposits, grouped under the name “Tamala Limestone”, have been investigated along thousands of kilometers of coastal Western Australia (WA). Relative-age diagenetic features of carbonate sand dunes or “eolianites” indicate that coastal ridges decrease in age seaward, reflecting coastal accretion during successive sea-level stands. Yellow- to red-stained quartz sands are associated with eolianites as pits, lenses, extensive beds, and even 40-m-high dunes.

A regional survey using whole-rock and land snail amino acid epimerization geochronology confirms the eolianite succession and provides a means of correlating widespread deposits along a steep climatic gradient and 16° of latitude. AMS ¹⁴C and uranium–thorium (U/Th) ages provide independent radiometric calibration of the amino acid ratios, and eolianite ages can be estimated using a parabolic kinetic model.

Over 90% of the sampled eolianite deposits comprise Aminozones A, C, E (125 ka), F?, G, and H, and correlate with interglacials from < 10 ka (Holocene) to ca. 500 ka. In contrast, at the type locality of the Tamala Limestone along the Zuytdorp Cliffs, the upper eolianite-paleosol units are characterized by advanced stages of cavernous weathering, pedogenesis, and recrystallization. In the same units, sediment and snail samples generally yield very low or non-detectable levels of amino acids. These factors along with the stratigraphic complexity of the deposits reflect an age much greater than a large majority of sites along the WA coastline. These findings encourage a revision of the existing classification and nomenclature of Quaternary carbonate deposits in WA, as well as a reexamination of the underlying mechanisms related to the formation and emplacement of both carbonate and quartz dunes.

© 2007 Elsevier Ltd. All rights reserved.

Keywords: Western Australia; Eolianite; Amino acid epimerization dating; Whole-rock; *Bothriembryon* land snails; Limestone diagenesis; Quaternary sea-level cycles; Tamala Limestone

1. Introduction

Coastal carbonate sediments and rocks representing subtidal, beach, and supratidal facies abound in Western Australia (WA), particularly from Shark Bay southward, where prevailing winds are generally onshore (Fig. 1). In Bermuda, which serves as a model for the interpretation of coastal eolianites, similar biogenic dune deposits were defined as “eolianites” by Sayles (1931). Sayles (1931) originally suggested that carbonate eolianites were glacial-age dunes deposits that migrate across exposed shelf areas.

*Corresponding author.

E-mail address: paulh@uow.edu.au (P.J. Hearty).

Bretz (1960) later reversed this perspective by demonstrating eolianites are a supratidal facies of coastal subtidal and beach deposits, and thus correlated them with highstand cycles which remains the prevalent global view. In WA, Fairbridge and Teichert (1953), and many others to follow, maintained a model of eolianite deposition during periods of glacially lowered sea level.

Eolianites generally form a broad coastal ridge parallel to the coast, and are volumetrically dominant on many subtropical coastlines (Brooke, 2001). However, with the exception of well-dated reef deposits ca. 125 ka (Stirling et al., 1995, 1998; O'Leary, 2007; O'Leary et al., in press, in review), there is little consensus in WA on virtually any aspect of carbonate dune deposits including their

Fig. 1. WA map showing study Regions I, II, and III. The steep climate gradient between cool temperate and dry tropical conditions encompassed by these regions are reflected in amino acid ratios. Image on right is an oblique satellite image of Shark Bay with site numbers: (1) Woomerangee Hill at Zuytdorp Cliffs, the type locality of the Tamala limestone; (2) Tamala Station; (3) Monkey Tarra; (4) Baba Head; (5) Eagle Bluff and Eagle Island; (6) Town of Denham; (7) Cape Peron (type locality of the Peron Sandstone); (8) Monkey Mia; (9) Hamelin Pool and stromatolites.

provenance, mechanism of transport (e.g., migration or aggradation), age of formation, and timing of dune emplacement relative to sea-level cycles. In this investigation, we focus on four important issues: (1) the abundance and distribution of Quaternary carbonate dunes (henceforth “carbonate dunes or eolianites”) of various ages; (2) the origin and significance of interbedded quartz deposits including dunes on the coastal plain; (3) the formation and age of the Tamala Limestone at the type locality along the Zuytdorp Cliffs, and; (4) the relevance and suitability of this type locality to represent eolianite deposits across WA (Table 1). We address these questions primarily through a regional chronostratigraphic survey of sites correlated and dated with amino acid epimerization (AAE) geochronology, AMS ^{14}C , and U/Th ages.

1.1. Eolianites in WA

The dearth of scientific information about eolianites in WA is partially attributed to the remoteness of the coastline outside of Perth, the sheer scale of the deposits, and to the limitations in available dating methods. The effective range of AMS ^{14}C is too short to be relevant to most Pleistocene deposits, while pristine *in situ* corals, required for U/Th dating, occur in only a handful of exposures of reef facies in the region. In contrast, eolianite ridges occur throughout coastal WA. The development of whole-rock (WR) aminostratigraphy (Hearty et al., 1992)

offers new opportunities to correlate and date Quaternary carbonates, particularly eolianites that generally lack macrofossils.

In Fig. 2, a facies model developed from Bahamian studies (Hearty and Kindler, 1997) represents eolianites as highstand supratidal facies. The highstand eolianite model has a strong basis in previous research. This investigation is preceded by a number of aminostratigraphic studies, which also employ coastal geomorphology, stratigraphy, and sedimentology from the Mediterranean (Hearty et al., 1986), Bermuda (Hearty et al., 1992; Hearty, 2002a), Bahamas (Hearty et al., 1996; Hearty and Kaufman, 2000), Hawaiian Islands (Hearty et al., 2000, 2005a, b; Hearty, 2002b; Fletcher et al., 2005), South Australia (Murray-Wallace et al., 2001), and WA (Murray-Wallace and Kimber, 1989; Hearty, 2003). These studies offer numerous examples of coastal facies relationships between shore and dunes, and demonstrate the effectiveness and reliability of the AAE approach. Through AAE, it is possible to group sites according to relative age, determine the relative abundance of sites of similar age, and to make generalizations about the likely correlation with several past sea-level cycles.

1.2. Interbedded deposits of quartz sand

In WA, eolianites consistently contain a small percentage of quartz and lithic grains, which are presumably

Table 1
Stratigraphic nomenclature and correlation of units in Perth and Carnarvon Basins, Western Australia

Stage	Teichert (1950)	Playford et al. (1975; 1976)	Playford (1997)	Logan et al. (1970; 1974)	Rottneest/ Perth area Murray-Wallace and Kimber, (1989) Hearty (2003) in bold	Shark Bay Eagle Bluff and Lookout (this study in Bold)	MIS correla- tion (this study)	
Holocene	Coastal limestone	Tamala limestone or eolianite (glacial lowstands only)	Holocene-recent interval	Modern/subrecent beach and dune	Unit x Eolianite and marine Unit m Marine quartz and carbonate	1		
Holocene beach and dune Unit VI								
Upper Pleistocene			Nilemah sands	paleosol	Unit l Quartz dunes	2 ↓ 4		
				Fish Hook Bay eolianites Unit V	Unit k Carbonate dunes	5a ↓ 5c		
				paleosol	Unit i-j Pisolith and quartz dune	5d		
Middle Pleistocene			Bibra Formation; marine reef; oolite; dunes	Rottneest Limestone/ Minim Cove - marine and eolianites Unit IV	Unit h Marine and eolianites	5e		
			Dampier Formation	Peppermint Grove marine and eolianites Unit III	Unit b-g Continental dune cycles; no marine?	7 ↓ 11-15		
				Carbla oolite?			Guilderton Quarry eolianite/ Pinnacles Unit II	Unit a Eagle Island? Baba Head; Tetradon yellow oolite
Major hiatus in coastal highstand record (MIS 15-25?)								
Lower Pleistocene			Tamala Eolianite	Yoganup Fm Unit I	Zuytdorp Cliffs	25?		
	Peron Sandstone	Old quartz dunes, Peron Peninsula; Monkey Mia	↓ >37?					

incorporated into carbonate beach and dunes during the erosive action of underlying continental sediments during marine transgressions (Hearty, 2003). The abundance of quartz sand is generally less than 10% in eolianites (Table 2), while soil and paleosol horizons generally contain higher proportions of quartz grains. In many examples, eolianites are interbedded with, or overlain by decimeters to meters of yellow to red, nearly pure (>95%) quartz sand (Allayialis, 2003). Pure quartz sands also infill pits in eolianite (Fig. 3).

The source of these quartz deposits is inextricably interwoven with the discussion of eolianites. One school of thought maintains that the quartz sand deposits result from dissolution of the carbonate sediments by meteoric waters, thus creating concentrations of quartz grains or “residual soils” (e.g., Bastian, 1996). An opposing view is that the quartz is distributed seaward (westward) across the coastal plain by migrating continental dunes during periods of lower sea level (or “desert expansion” of Semeniuk and Glassford, 1988).

Shallowing-upward sequence on sedimentary shorelines

Fig. 2. A subtidal to supratidal (dune) structural transect across a typical sediment-dominated coastline (after Hearty and Kindler, 1997). The shallowing-upward facies model is representative of typical Holocene and older interglacial sequences of Western Australia, and implies association with sea-level highstand intervals.

1.3. The type locality of the “Tamala Limestone”

The Zuytdorp Cliffs of central coastal WA maintain an important distinction as the type locality of the Tamala Limestone, representing Quaternary coastal eolianites in the Shark Bay area (Logan et al., 1970) and across all of WA (Playford et al., 1975, 1976). Playford's (1990, 1997) broader definitions would encompass widespread and voluminous carbonate marine and eolianite deposits along 5000 km of WA coastline and span over 1 Ma. Despite this impressive status, neither a detailed description of the deposits nor any geochronology has been published from the type area.

Teichert (1950) first referred to the assemblage of marine and eolianite deposits along the coast of WA as the “coastal limestone” (Table 1). Brian Logan and colleagues later discussed (in Gill, 1968) and defined (Logan et al., 1970, p. 52) the “Tamala eolianite (as) a vast complex of dunes which rise...up to 500 ft above present sea level”. They specified no type locality, but presumably were referring to the adjacent Zuytdorp Cliffs several kilometers west of Tamala Cattle Station (Fig. 1). They characterized the deposits as “consolidated to unconsolidated, coarse- to medium-grained, skeletal-fragment grainstone which commonly shows large-scale (eolian) cross-bedding”. They

assigned an early Pleistocene age to these deposits (their Table 2, p. 51). The name “Tamala” was thus adopted to represent Quaternary eolianites, first in Shark Bay, and subsequently in all of WA (Fig. 1).

Playford et al. (1975) identified a site at +258 m elevation at Woomerangee Hill (26.884°S. lat., 113.749°E. long) along the Zuytdorp Cliffs as the type locality (Fig. 1), and measured a 251 m section. Playford et al. (1975) concurred with Logan et al.'s (1970) general description of the sediments, but regarding the age, they stated (p. 307) “the eolianite extends below present sea level, and it seems more likely that it formed during one of the glacial regressions, probably in late Pleistocene times”. Playford et al. (1976) then expanded the definition to include carbonate dunes in the Perth and Carnarvon Basins, and reiterated their opinion that the deposits were formed during a late Pleistocene glacial age. Playford (1990, p. 20) later broadened his definition to include “accumulation during (multiple) glacial periods.” And more recently, Playford (1997, p. 794) stated “the Tamala Limestone accumulated as coastal sand dunes, mostly during periods of lowered sea level during the Pleistocene and early Holocene, when wide areas of the continental shelf were exposed, and carbonate productivity was high.”

Table 2

Carbonate content and color of quartz pits, lenses, beds and dune deposits from sampled localities in WA (from Allayialis, 2003)

Site location	Sample ID	Dated sample (this study)	Estimated age of quartz (MIS) ^a	% Carbonate	Munsell (2002) moist color
Port Smith	BSM2x	BSM1ab	4–2	9	5YR 5.5/6
Pardoo	BPO1	Continental dune	Post 5?	0	10R 5/8
Vlaming Head	NVL1;		Post 5e	12	2.5YR 4/8
	NVL1d		Pre 1	9	
Exmouth	WEX1a				
	WEX2a		Post 9/11	7	5-7.5YR 6/6
Vlaming Head	WVL3a			8	5YR 6/6
Milnya	MY1	Continental dune		0	2.5YR 5/8
Broadhurst North	SBN1a;			0	2/5YR 4.5/8
	SBN1e			0	2.5YR 4/8
Billabong	BB25	Continental dune		0	5YR 4.5/6
White Point	WWP2b	WWP2a 0.42	Post 5e	0	10YR 7/8
		WWP1a 0.40			
Leeman	WLE2b	WLE1a 0.35	Post 5e	1	7.5YR 5.5/6
		WLE2x 0.22	Pre early 1		
Jurien	WJR1b	WJR1a 0.46	6 or 4	0	7.5YR 5/8
Jurien	WJR2c	WJR2a 0.49	6 or 4	0	10YR 7/6–8
Cervantes	WCP2e;	WCP1x;3x	Post 11	0	10YR 7/8
	WCP4e;		Pre 1–4–2	0	10YR 7/8
	WCP5e			38	10YR 7.5/6–8
Lancelin	WLN1a	Continental dune	4–2	0	10YR 7/6–8
Lancelin	WLN2b	WLN2a	4–2		10YR 6.5/8
Lancelin	WLN3a,	WLN3a	Post 11?	55	10YR 7.5/4
	WLN3e	WLN3e	Pre 1	0	10YR 7/8
Kalgoorlei	KAL1	Continental dune		0	10YR 6.5/8
Guilderton	PGU1b	PGU1a	Post 11	0	10YR 6.5/6
Mt Eliza	PEZ3b	PEZ1a	Post 11–13	2	10YR 7/8
Fairbridge Bluff, Rottne I (Rotto)	RFB1b;	RFB1a	Post 5e	13	10YR 8–7/4
	RFB1d		Pre 1		10YR 7.5/3–4
Mary Cove, Rotto	RMB2d;		Post 5e–6?	38	10YR 6/3
	RMC1b				10YR 7/3
Mimim Cove, Rotto	PMC1e	PMC1ad	Post 5e	0	7.5YR 6.5/6–8
Jeannie's, Rotto	RJL2b	RJL2a 0.37	Post 5e	45	7.5YR 6.5/4
Mandurah Q	WMQ1a	Continental dune	4–2	0	10YR 7.5/4
Tims Thicket	PTT2d	PTT1a	Post 5e	4	10YR 8/6
Mandura Q	WMA1	Continental dune	Post 11	0	10YR 8/8
Esperance (Esp) Q	EEQ1b	EEQ1a	4–2	0	2.5Y 8/2/3??
Twilight Bay, Esp	ETW1b	ETW1a	Post 5e	0	2.5Y 7.6?
Nine mile Bch, Esp	ENI5c	ENI5a/b	Post 5	0	10YR 8–7.5/6
		ENI5d	Pre 1		
Nine mile Bch, Esp	ENI3b	ENI3a	Post 5e–Pre 1	24	10YR 6/4
Foul Bay	MFU2e	MFU2	Post 5e	5	7.5YR 7/4
Cozy Corner	CCR1d	CCR1c	Post 5e–Pre 1	0	10YR 5.5/6
QB Quarry	QBY2d	QBY2b	Post 11–Pre 1	5	7.5YR 5.5/6
Blossom Bch	ABB2a	ABB2a	Pre 7/9	0	7.5YR 6–5/8
Ankatel N Perth	WAN1b(1)	Continental dune	4–2	0	2.5Y 7/2–3
	WAN1b(2)	Continental dune	4–2	0	2.5Y 7/2–3

^aMarine isotope stage (MIS) correlation of quartz deposits are based on AAR determinations from carbonate deposits in this study.

2. Geologic and climatic setting

2.1. Field area and climatic setting

The study area extends over several thousand kilometers of coastal WA from Esperance to Broome (Fig. 1) covering over 16° (34–18° South) of latitude. Climate ranges south to north from cool temperate to tropical, and seasonally wet to extremely hot, windy, and

dry. The mean annual temperature varies from about 17 °C (mean minimum and maximum annual 14–19.7 °C) at Cape Leeuwin to 27 °C (mean minimum and maximum annual 21–33 °C) at Cape Range, while rainfall across the same gradient ranges from 600–1000 to 100–200 mm/yr (Australia Bureau of Meteorology: www.bom.gov.au). Evaporation rates from south to north are about 1200 mm/yr at Cape Leeuwin to over 3000 mm/yr at Cape Range.

Fig. 3. Pure quartz (“Q”) deposits in pits in carbonate dunes (“C”) near Lancelin (WLN2 (A)), lenses and pits at Esperance quarry (EEQ1 (B)), extensive meter-thick beds at Nine Mile Beach in Esperance (ENI3b (C)), and 30-m-high quartz dunes at Mandurah Quarry (WMQ1 (D)), which produced Aminozone *G/H* values on eolianite at its base. Large quartz dunes (like (D)) of middle and late Pleistocene age (Table 2) are found at many localities along the coastal plain of SW WA.

The abundance of Quaternary eolianites in WA is largely dictated by high shelf productivity (warm waters: Leeuwin Current), general aridity, and persistent and strong prevailing onshore winds. A global inventory of eolianites (Brooke, 2001) indicates that their greatest abundance occurs between 30° and 40° north and south latitude. South of 30° in WA, the climate is seasonal and strongly influenced by deep low-pressure fronts sweeping across the Southern Ocean, with strong prevailing westerly winds, and heavy onshore ocean swell. North of about 25°S, the patterns reverse with dry easterly continental winds flowing offshore of the Great Sandy Desert creating extremely dry and frequently dusty conditions and fewer eolianites. These continental wind patterns appear to contribute to the

sedimentology with greater percentages of quartz and reddened sediments appearing the coastal eolianites.

2.2. Field stratigraphy and geomorphology

The Quaternary stratigraphy of WA generally consists of a succession of bioclastic dune and shoreline facies intercalated with or capped by either weak interglacial soils called protosols (Vacher and Hearty, 1989), or yellow to red clay and/or quartz-rich soils or paleosols (i.e., buried soils) (Fig. 4). In the Bahama Islands, Hearty and Kaufman (2000) used the term “couplets” to describe the eolianite-soil pairs, and correlated the former with interglacial highstands, and the latter with lowstand events,

Fig. 4. Montage of typical late and middle Pleistocene eolianites from WA: (A) Succession of late Pleistocene eolianites and weak paleosols at Fish Hook Bay, Rottneest Island (arrow: person for scale). (B) Holocene dunes abutting early Pleistocene eolianites at Conto's Beach south of Margaret River. (C) Late Pleistocene foreset beds at Hamelin Bay south of Margaret River. (D) Middle Pleistocene eolianites in a small quarry above Quarry Bay, Cape Leeuwin. (E) Quartz dunes and "pinnacles" at Nambung National Park. (F) Pinnacles emerging from middle Pleistocene eolianite 5 km east of Guilderton.

when the source shoreline shifted far seaward from the modern coastline.

Most of the WA eolianites consist of weakly to moderately cemented, fine-to-coarse carbonate sand displaying large-scale festoon and planar cross-bedding, generally on windward and leeward flanks, respectively (Fig. 2). Microscopic examination reveals the primary bioclastic components in eolianites (and WR AAE samples) to be mollusks, calcareous red algae, foraminifera, and echinoderms (Kendrick et al., 1991; Kinna, 2002). Most taxa are warm-water species with a similar biogeographic distribution to modern highstand sediment (James et al., 1999), with no biogeographic suggestion of colder, glacial-stage oceanographic conditions on the shelf.

With the exception of a reach of coastline SW of Broome (Hearty et al., 2006), no ooids or oolites are documented from open ocean coastlines of WA (James et al., 2004). However, both Pleistocene and Holocene ooids are abundant in the hypersaline waters near the stromatolite

formations (Logan et al., 1970, 1974) deep within Shark Bay.

The primary biogenic grain textures and the original colors of the skeletal organisms are progressively lost with greater age of the limestone, while the progression of CaCO_3 recrystallization (aragonite $\rightarrow$ high-Mg calcite $\rightarrow$ low-Mg calcite), secondary mineral precipitation (increasing rock density and hardness), and dissolution and cavernous weathering increase with age (Kindler and Hearty, 1996; Hearty et al., 1996, 2005a; Hearty, 1998). The amino acid concentration is progressively decreased with advancing age and state of diagenesis of the carbonate medium (Roof, 1997; this study). Soils capping limestone deposits increase in richness, hue and chroma (yellows and browns to reddish brown and red) with greater age (Hearty and Kindler, 1997; Hearty, 1998; Hearty et al., 2005a). Diagenetic "grades" I (unconsolidated carbonate sand) to V (recrystallized limestone), defined by Land et al. (1967) for a succession of Bermuda limestones, are appended here as they relate to the Quaternary of WA (Fig. 5).

3. Approach and methods

3.1. Overview

To address many of the observations and issues presented above, an extensive survey of coastal eolianites and quartz deposits was undertaken in WA since 2001. Sites were examined and samples collected, and various analyses were made to determine the composition and

possible ages of the deposits. The findings of a regional aminostratigraphic investigation are summarized here, and comparisons are made with other tectonically stable carbonate-producing Quaternary shoreline and AAE sequences of the world.

At each site, stratigraphic sections were logged and photographed. Site locations were assigned names based on the nearest topographic landmark on either 1:100 000 or 1:250 000 map sheets. Samples from over 200 sites were examined and analyzed during our field investigations. These sites are too numerous for plotting in Fig. 1, but GPS coordinates are provided for most sites (Table A1). Those few sites lacking GPS coordinates can be located by place names on topographic sheets of the region. With the exception of the widespread Holocene and modern dunes, nearly all indurated eolianite outcrops encountered along most road cuttings and 4-wheel-drive accessible coastal exposures were described and sampled. We thus consider our collection of sites to be a reasonably comprehensive and unbiased survey of Quaternary exposures in the region. Included among these sites are two localities along the Zuytdorp Cliffs, including Playford's type section at Woomerangee Hill and another north of Tamala station called "Monkey Tarra".

Fig. 5. Mineralogical, thin-section, and relative-age characteristics (rock hardness and Munsell (2002) soil color) defining diagenetic grades from a succession of Holocene (grade I) to early Pleistocene (grade V) carbonates from Bermuda (Land et al., 1967; this study). A large majority of eolianite outcrops along the WA coastline are diagenetic grades I–III. Grades IV and V characteristics are rarely observed along the coastal plain.

3.2. Atmospheric and marine reservoir effects on ^{14}C ages

AMS ^{14}C ages were determined on fossil *Bothriembryon* shells and carbonate sediment samples (Table 3) at the Lucas Heights facility near Sydney, Australia. Both sample materials were prepared in the same manner as AAE samples (below). Radiocarbon ages were calculated based on percent modern carbon with reference to $\delta^{13}\text{C}$ normalized to

Table 3
AMS ^{14}C ages and amino acid ratios of samples analyzed for *Bothriembryon* land snails, whole rock and charcoal

ANSTO lab ID	Field ID	$\delta^{13}\text{C}$ (‰)	^{14}C age (BP)	$\pm 1\sigma$	Calibrated age (cal BP)	$\pm 2\sigma$	A/I mean	A/I $\pm 1\sigma$	A/I N
<i>Bothriembryon land snails</i>									
OZG917	STL3	−10	Mod	Mod	Modern		0.02	0.00	2
OZG459	SMT1(DIO)	−7.4	1715	40	1563	138	0.05	0.02	3
OZG460	SMT4(SIO)	−6.9	5645	50	6370	94	0.17	0.06	3
OZG461	SEB4I	−6.6	9380	60	10,528	200	0.27	0.04	4
OZG462	SBS1as	−3.5	3620	50	3860	183	0.17	0.06	5
OZG916	RCY1b	−6.8	31,500	280	Non-finite		0.33	0.02	2
OZG471	SLL1a	−5.2	55,500	5600	Non-finite		0.57	0.01	1
<i>Carbonate-rich sediment (cements removed)</i>									
OZG463	RFB1z	0.1	2015	40	1581	112	0.05	0.00	2
OZG464	RAJ1a	0.5	9270	70	10,093	190	0.18	0.02	2
OZG465	BSM1a(1)	2.6	3515	45	3401	118	0.10	0.02	2
OZG466	RSP1e	0.2	22,960	160	Non-finite		0.22	0.01	4
OZG467	RHQ1a(2)	0.2	19,070	150	22,234	333	0.23	0.02	2
OZG468	SMT1(DIO)	−0.2	19,100	140	22,262	263	0.42	0.04	2
OZG469	SMT4(DIO)	−0.5	32,240	260	Non-finite		0.72	0.02	2
<i>Charcoal</i>									
OZG456	WAN1b(1)	−27.8	3430	50	3615	176	–	–	–

Note: Ages were calibrated to calendar years using CALIB v. 5.0.2 and a standard marine reservoir correction for the skeletal sediment (Reimer et al., 2004; Stuiver et al., 1998, 2005). Calibrated ages are reported as median probability and $\pm$ one half of the 2σ age range.

Table 4
U/Th ages on coral from Western Australia sites confirming a correlation with MIS 5e

Site	Location	U (ppm)	Th (ppt)	$\delta^{234}\text{U}^a$	$\pm 2\sigma$	$^{230}\text{Th}/^{238}\text{U}^b$	$\pm 2\sigma$	$^{232}\text{Th}/^{230}\text{Th}$	Age (ka) ^c	$\pm 2\sigma$	Initial $\delta^{234}\text{U}^d$	$\pm 2\sigma$	Ref ^e
<i>Margaret River</i>													
MFU1a	Foul Bay	3.19	36.96	118.58	0.83	0.7124	0.0020	3212	107.42	0.42	160.79	1.05	1
CMP1	Cowaramup	3.10	38.30	108.38	0.47	0.7591	0.0033	4380	122.05	0.97	153.19	0.74	1
<i>Cape Burney</i>													
GCB2	Cape Burney	2.76	34.52	104.63	1.60	0.7738	0.0036	52810	127.30	1.10	150.10	2.10	1
<i>Shark Bay</i>													
SBA1	Baba Head	2.81	34.79	107.07	1.10	0.7601	0.0018	3427	122.64	0.56	151.59	1.47	2
STG5	Tetradon Loop	2.88	35.94	103.81	1.30	0.7720	0.0032	14,766	127.00	1.00	148.80	1.75	2
STG1	Tetradon Loop	2.46	31.09	106.57	0.66	0.7801	0.0030	72,614	128.80	0.92	153.50	0.96	2
STG4	Tetradon Loop	1.99	24.70	105.73	1.14	0.7683	0.0025	39,536	125.40	0.76	150.80	1.50	2
SPI4	Pelican Island	2.92	35.93	107.96	1.05	0.7612	0.0022	23,948	122.80	0.65	152.90	1.40	2
SPI5	Pelican Island	2.36	28.64	107.80	0.88	0.7500	0.0019	9177	119.60	0.58	151.30	1.15	2
SGN1	Gladstone N	2.76	35.47	113.75	1.05	0.7942	0.0023	90,678	131.30	0.79	165.20	1.40	2
SGN3	Gladstone N	3.25	41.08	116.18	0.72	0.7816	0.0015	18,740	126.87	0.47	166.40	0.96	2
SMM3a	Monkey Mia	2.28	28.43	109.06	0.97	0.7717	0.0024	5347	125.60	0.75	155.70	1.30	2
SMM4	Monkey Mia	3.79	46.69	106.04	0.57	0.7603	0.0016	61	122.94	0.46	150.20	0.77	2
<i>Cape Cuvier</i>													
LCC9a	Cape Cuvier	1.24	15.67	108.68	0.70	0.7787	0.0022	4938	127.80	0.69	156.13	0.93	3
LCC9b	Cape Cuvier	2.93	36.75	107.97	0.93	0.7748	0.0039	10,949	126.80	1.22	154.68	1.31	3
LCC9c	Cape Cuvier	1.43	18.40	107.44	0.54	0.7948	0.0040	10,814	133.15	1.25	156.71	0.85	3
LCC9d	Cape Cuvier	2.67	33.17	104.93	2.40	0.7672	0.0080	6282	125.24	2.41	149.66	3.20	3
<i>Cape Range-Ningaloo</i>													
NJZ1	Jacobesz	2.02	25.26	108.60	1.22	0.7675	0.0022	929	124.47	0.70	154.55	1.580	3,4
NJZ2	Jacobesz	2.14	26.58	108.52	0.81	0.7653	0.0020	6291	123.84	0.61	154.16	1.040	3,4
NYC1a	Yardie Creek	2.32	28.94	103.67	1.55	0.7678	0.0021	8483	125.74	0.76	148.07	2.020	1
NYC1b	Yardie Creek	2.45	33.59	111.65	1.19	0.8433	0.0020	22	148.48	0.84	170.09	1.640	1

Note: Additional ages from several calibration sites are available in Stirling et al. (1995, 1998). Ages in bold represent strictly reliable ages (see text).

^a $\delta^{234}\text{U} = \{[(^{234}\text{U}/^{238}\text{U})/(^{234}\text{U}/^{238}\text{U})_{\text{eq}}] - 1\} \times 10^3$. $(^{234}\text{U}/^{238}\text{U})_{\text{eq}}$ is the atomic ratio at secular equilibrium and is equal to $\lambda_{238}/\lambda_{234} = 5.4891 \times 10^{-5}$, where λ_{238} and λ_{234} are the decay constants for ^{238}U and ^{234}U , respectively, adopting half-lives of Cheng et al. (2000).

^b $[^{230}\text{Th}/^{238}\text{U}]_{\text{act}} = (^{230}\text{Th}/^{238}\text{U})/(\lambda_{238}/\lambda_{230})$.

^cU/Th; Table 4 ages are calculated iteratively using: $1 - [^{230}\text{Th}/^{238}\text{U}]_{\text{act}} = \exp\left(\frac{\lambda_{230}}{\lambda_{238}} - (\delta^{234}\text{U}(0)/1000)(\lambda_{230}/(\lambda_{230} - \lambda_{234}))(1 - \exp(\lambda_{234} - \lambda_{230}T))\right)$ where T is the age in years and λ_{230} is the decay constant for ^{230}Th . $\lambda_{238} = 1.551 \times 10^{-10} \text{ yr}^{-1}$; $\lambda_{234} = 2.826 \times 10^{-6} \text{ yr}^{-1}$; $\lambda_{230} = 9.158 \times 10^{-6} \text{ yr}^{-1}$. Strictly reliable ages have $^{234}\text{U}_{\text{initial}} 146.6 \pm 5\%$.

^dThe initial value is given by $\delta^{234}\text{U}_i = \delta^{234}\text{U}(0)\exp(\lambda_{234}T)$, where T is the age in years.

^eReferences: 1 = this study; 2 = O'Leary et al. (in press) Shark Bay; 3 = O'Leary et al. (in review) Cape Cuvier; 4 = Hearty et al. (2007).

–25‰. As dune sediments are composed of skeletons of organisms formed in shallow marine waters on the shelf, a marine reservoir age was applied to all AMS ^{14}C ages using CALIB 5.0.2 (Reimer et al., 2004; Stuiver et al., 1998, 2005). We chose the standard reservoir age because, although the evidence from the region is sparse, the ages are not significantly different from 400 yr (calib.qub.ac.uk/marine/index). All ^{14}C ages were calibrated to calendar years before 1950 AD (cal BP) and reported as median of the probability distribution of the calibrated ages with errors of one-half of the 2σ calibrated age range.

We recognize that land snails may carry with them a considerable dead carbon anomaly (100 s to 1000 s of year) inherited from grazing on radiocarbon-dead limestone (Goodfriend and Stipp, 1983; Goodfriend and Gould, 1996). It is thus possible that several of our shell ages may be somewhat too old, but how much older is difficult or impossible to ascertain. As such, we do not apply a dead-carbon correction. It is encouraging that one live-collected

shell sample from Shark Bay (OZG917) was tested and returned a modern AMS ^{14}C age.

3.3. U/Th methods

U/Th measurements on corals were performed using a Neptune MC-ICPMS at the Research School of Earth Sciences, Australian National University. Measurements were conducted using a combination of simultaneous multiple-Faraday cup and ion counter protocols. Following the procedures of earlier workers (e.g., Stirling et al., 1995, 1998; McCulloch and Esat, 2000), coral ages (Table 4) were screened for potential U and Th loss or gain based on the following criteria:

1. We consider the calculated $\delta^{234}\text{U}_{\text{initial}}$ to be the best quantitative test for open-system behavior in corals. For a coral age to be considered strictly reliable $\delta^{234}\text{U}_{\text{initial}}$ values should reflect a modern seawater value of $146.6 \pm 4\%$.

2. The total uranium concentration of fossil corals should approximate modern coral values of about 3 ± 0.5 ppm of uranium.
3. Fossil corals should be free of allochthonous ^{230}Th , as indicated by the absence of detrital ^{232}Th (< 1 ppb).
4. Corals should show primary aragonitic structures or have $< 2\%$ calcite concentration.

Primarily strictly reliable coral ages are reported in Table 4; however, several other samples of particular interest with less-reliable ages are also presented. We emphasize that this is not intended to be a study of U/Th techniques. We refer to these ages only to confirm that the deposits equate with MIS 5e, and then to provide calibration for mean AAE data from these sites. More detailed investigations of aspects of U/Th geochronology from numerous WA sites are available in Stirling et al. (1995, 1998), Hearty et al. (2007), O'Leary et al. (in press, in review), and several forthcoming papers.

3.4. Amino acid geochronology

A total of ~500 samples were analyzed for AAE from three latitudinally defined regions (Fig. 1): Region I—Cape Leeuwin to near Geraldton ($34\text{--}29^\circ\text{S}$ lat.); Region II—Geraldton to Shark Bay and Cape Cuvier ($29\text{--}25^\circ\text{S}$ lat.), and Region III—Cape Cuvier to Cape Range and Port Smith near Broome ($25\text{--}18^\circ\text{S}$ lat.). Sedimentological analyses of many of the samples were conducted by Honours and Ph.D. students from the James Cook University (Townsville, Queensland) including Kinna (2002; biogenic carbonates), Allayialis (2003; quartz deposits), Donald (2003; Shark Bay stratigraphy and sedimentology), and O'Leary (2007 and following papers; WA geochronology).

A large database of AAE analysis of WR and the ubiquitous WA land snail *Bothriembryon* provides the primary device for site-age comparisons across WA. Additional details of the WR sample preparation procedures and protocols are presented in Hearty and Kaufman (2000). Briefly, WR samples were milled and sieved, split into two equal samples, and each prepared and analyzed to assess the analytical uncertainty. Samples were gently disaggregated and sieved to obtain the 250–850 μm sand fraction. Previous studies have shown that this size range is typical of beach and dune facies, and tends to exclude fine intergranular cements as well as large grains or shells that could disproportionately influence the sample result. The A/I value is a measure of the extent of *epimerization* of the common amino acid isoleucine. In living organisms, the A/I value after preparation is initially near zero (~ 0.015) and increases with age to a ratio of ~ 1.30 . Several modal classes or “aminozones” of D-alloisoleucine/L-isoleucine (or A/I) ratios from both sample materials are calibrated with numerical ages by AMS ^{14}C and U/Th ages from this and other studies. Changing thermal histories across the WA climatic gradient from Cape Leeuwin to Cape Range yield different values, such that mean A/I for

Aminozone E in Region I (~ 0.30) will be lower than Regions II and III (~ 0.50 to ~ 0.56). Correlation of the Aminozone E across the gradient is accomplished by independent U/Th ages on corals. All AAE analyses were performed at the Amino Acid Laboratory of Northern Arizona University (D. Kaufman, Director) following the standard protocols of Miller and Brigham-Grette (1989).

As the WR method analyzes aggregates of skeletal grains that form over time, entirely “modern” ratios are not expected. For example, modern beach sand from near Fairbridge Bluff on Rottnest Island yields an AMS ^{14}C age of 1581 ± 112 cal BP and a mean A/I of 0.049 ± 0.000 (2) (Table 4 and Table A1). This sample presumably includes a mix of sand grains from late Holocene to modern age. Mixing of intraclasts from older interglacial deposits may occur in some cases, but as the concentration of amino acids decreases exponentially with age, limited percentages of reworked grains do not yield sufficient concentration of amino acids to significantly alter the A/I value (Hearty and Kaufman, 2000). In contrast, however, the mixing of carbon-dead grains into younger deposits has a profound effect on ^{14}C ages, which can be independently assessed by AAE as will be demonstrated below.

Three to eight *Bothriembryon* shells were analyzed and averaged from a total of about 40 stratigraphic levels. *Bothriembryon* are abundant in most stratigraphic units around Shark Bay, and more sparsely in the adjacent Zuytdorp Cliffs. In all cases, the shells were sub-sampled from the same area of the apertural lip. This taxon demonstrates moderately variable AAE results compared to other land snails, possibly due to the more open microstructure of the fossil shells.

3.5. Amino acid diagenesis: loss of amino acids with age

Previous studies have determined that amino acid concentration decreases exponentially with greater sample age (Hearty and Kaufman, 2000; Hearty et al., 2004a). If fossil or WR sediment samples are generally well preserved, and there is no evidence for *unusual* diagenesis (such as associated cavernous weathering, extreme Fe staining, complete recrystallization), then there is a threshold age at which samples yield insufficient levels of amino acids to be accurately measured. “Insufficient levels” translates to detection at the level of hundreds of picomoles with the ion exchange technique. A demonstrated effect of diagenesis and leaching is the selective loss of D-alloisoleucine (“A” of A/I), relative to L-isoleucine, resulting in decreasing A/I in the most altered samples (Roof, 1997). For this reason the maximum A/I values of a set of several from a single stratigraphic level are considered to be the most reliable indication of the deposit age. The terminal condition of the samples is the complete loss of amino acids (non-detectable on the HPLC), typically in the oldest recrystallized samples. As will be examined below, we suggest this progression is time dependent, and can thus be used as a relative-age indicator within restricted climate regimes.

4. Results

4.1. Regional aminostratigraphy of coastal eolianites and marine deposits in WA

The primary objective of this study is to provide an independent assessment of the frequency of occurrence (the number of sites yielding similar ages) and the range of ages of eolianite deposits along the coastal zone of WA. This study does not attempt to provide precise age estimates of individual

sites, but only to draw reasonable generalizations on deposit ages (i.e., Holocene, late, middle, or early Pleistocene with the latter two each comprising several interglacials) from WR and land snail *A/I* data across the three regions delineated in Fig. 1.

4.2. Aminozones A–H from WR and *Bothriembryon* samples

WR histograms defined from Regions I–III show the frequency and distribution of mean *A/I* values for both

Fig. 6. Histograms of whole-rock (WR) and *Bothriembryon* *A/I* values in SW West Australia south of 29° latitude (Region I), central 29–25° latitude (Region II), and NW 25–18° latitude (Region III). The influence of warmer climate history to the north is indicated by higher calibrated mean WR *A/I* from U/Th calibration sites (arrows) across the region (Table 4; Stirling et al., 1995, 1998). Similar regional patterns are revealed in dunes and soils containing *Bothriembryon* land snails.

Table 5
Aminostratigraphy of Regions I and II of Western Australia (WA) determined in this study

Amino-zone	Region I: Perth (P) Rottneest (R)	Representative locality SW WA	Region II Shark Bay and Zuytdorp Cliffs	Representative locality (Shark Bay area)	MIS correlation
A3	0.05(1) 0.11 ± 0.02(15)	R: Salmon Bay R: Army Jetty; Herschell Q	0.06 ± 0.01(3) 0.28 ± 0.02(6)	Hamelin Pool Unit l, Eagle Lookout,	Modern to mid 1
A1/C? C/D	0.21 ± 0.03(8) 0.23 ± 0.03(74)	R: South Point R: South Point and Fish Hook Bay	0.41 ± 0.01(2)	Goulet Bluff Unit k, Eagle Lookout	Early 1 or reworked C 5a/5c
E2 ~125 ka E1	0.32 ± 0.04(42)	P: Minim Cove R: Fairbridge Bluff	0.50 ± 0.02(8) 0.62 ± 0.03(5)	Baba Head Gladstone N.	5e
F?/G	0.41 ± 0.01(4)	P: Peppermint Grove	0.67 ± 0.01(3)		7/9?
H	0.49 ± 0.04(6) 0.52 ± 0.02(2)	P: Fremantle Fort Guilderton Q.	0.97 ± 0.01(2)	Baba Head yellow oolite	11
	0.72 ± 0.02(1)	Cape Leeuwin: Quarry Bay high eolianite in Q			
BRS				Base of Cape Peron sections?	12–24?
K	> 780 ka ND	Leeuwin-Naturaliste ridge	Low A/I to ND	Zuytdorp Cliffs	25–37?

ND = non-detectable. BRS = Big red soil (see text). Numerical values are D-alloisoleucine/L-isoleucine (*A/I*) values (mean, $\pm 1\sigma$, (*N* = number of samples or sites)) from whole rock limestone samples. Ratios from equal-age sites increase with warmer climate history: SW WA coolest and NW WA warmest.

Fig. 7. Gradient of MIS 5e whole rock *A/I* values from U/Th-dated coral sites across coastal Western Australia. U/Th ages are from Table 4 and Stirling et al. (1995, 1998).

sample materials from 200+ sites (Table A1; Fig. 6). The Aminozones A–H nomenclature, indicated on the tables and histograms, represents modal classes of individual *A/I* values of increasing age (Table 5), as defined in previously mentioned studies. Aminozone A is correlated with the Holocene (MIS 1) deposits, and verified by ^{14}C analyses (Table 3). Aminozones C and E equate with the last interglaciation and encompass MIS 5a through 5e. U/Th ages on corals from cited studies and Table 4 confirm the correlation of Aminozone E with MIS 5e and provide a calibration for *A/I* means from each region (arrows in Fig. 6). Calibrated MIS 5e (125 ka) *A/I* values (Fig. 7) increase regionally between Margaret River and Cape Range between latitude 34° and 20° South ($r^2 = 0.86$).

Older Aminozones F?, G, and H probably correlate with middle Pleistocene interglacials and marine isotope stages (MIS) 7? to 11 (“F?” and “7?” are queried as we suspect coastal deposits of this age are rarely found above sea level

on stable coastlines, as discussed in Hearty, 2002a). As expected, the number of sites associated with older Aminozones F? through H are fewer compared with Aminozones A–E; older deposits are eroded or buried during younger coastal sedimentary events.

WR samples appear to provide regionally consistent *A/I* data for deposits up to Aminozone H. However, because of the slowing epimerization rate and lower amino acid concentrations, some older middle Pleistocene correlations may be off by plus or minus one interglacial. Beyond Aminozone H, older deposits commonly yield lowered *A/I* or non-detectable levels of amino acids. For example, a massive eolianite at Mt. Eliza (PEZ1) in King’s Park, central Perth, yielded trace to non-detectable levels of amino acids in three of four samples, averaging only $\sim 11.7\text{ nM/mg}$ (*A* + *I*). A measurable but unreliable *A/I* of 0.188 was determined from the remaining sample with a concentration of only 17.3 nM/mg. Mt. Eliza lies about 3 km landward of the Peppermint Grove site, which is correlated with MIS 7 or 9 (Murray-Wallace and Kimber, 1989; Hearty, 2003). Based on the 2–4 m higher-than-present sea level indicated by the subtidal–intertidal facies in the deposit, we suggest MIS 9 is the more likely correlation. A thermoluminescence age of >422 ka (Price et al., 2001) confirms that Mt. Eliza is certainly older than Peppermint Grove and most likely much older than Aminozone H (~ 400 ka).

4.3. Age estimates on WR carbonate sediment and shells using apparent parabolic kinetics (APK)

Epimerization kinetics for WR samples generally conforms more closely to a first-order reversible reaction than

for mollusks (Hearty et al., 1992). Nonetheless, APK can be used to estimate WR deposit ages. As mentioned, U/Th calibration at ca. 125 ± 5 ka of Aminozone E A/I values is provided at numerous sites along the WA coastline (Table A1 and Table 4; Fig. 6). Holocene deposits are generally <10 ka, as determined with AMS ^{14}C (Table 3) and low A/I values. The estimated ages of intermediate aminozones (D, C, and B) may be interpolated on the basis of an APK model (Mitterer and Kriausakul, 1989). Ages of aminozones older than E (125 ka) must be extrapolated as few middle or early Pleistocene calibrations are generally available in WA.

APK age estimates, based on A/I at 125 ka *per region* are based on the equation:

$$t = [(A/I)_s / m_c]^2, \quad (1)$$

where m_c is the calibrated rate of AAE = $[(A/I)_c / t_c^{1/2}]$, $(A/I)_s$ the A/I of the sample, and t the time in years.

MIS 5e calibrations (Table 4; Table A1) of WR A/I at 125 ka are: Region I at 0.32 ± 0.04 (8) (Foul Bay, Fairbridge Bluff, and Leander Point); Region II at 0.50 ± 0.03 (8) (C. Burney and Baba Head); and, Region III at 0.56 ± 0.07 (4) (Jacobs Rd). In all cases, WR A/I were determined from the marine sediment matrix enclosing or immediately adjacent to corals used for calibration. From APK, the maximum ages estimated from the highest and most reliable A/I s from Regions I, II, and III consistently range between 300 and 500 ka, correlating with several interglacials between MIS 9 through 13 (Table A1; Fig. 6).

4.4. APK on *Bothriembryon* land snails

Land snails contained in dunes immediately overlying MIS 5e marine deposits are assumed to date to latest MIS 5e, and therefore are assigned a slightly younger calibration age of 115 ka is assumed. *Bothriembryon* are only abundant in Regions I and II, where calibration A/I s are 0.34 ± 0.07 (4) (Foul Bay, Margaret River) and 0.83 ± 0.03 (20) (Eagle Bluff and Lookout, Shark Bay; Fig. 8A), respectively. In previous studies (e.g., *Poecilozonites* from Bermuda; Hearty et al., 1992, 2004b), empimerization kinetics in land snails generally exhibit parabolic pathways, compared to WR, and appear to do so for *Bothriembryon*, up to the last interglacial when independent calibration is considered over the past 130 ka (Fig. 8B). The estimated ages of aminozones based on *Bothriembryon* older than Aminozone E calculated by APK suggest a range of ages up to about 600 ka.

The stratigraphy of Shark Bay is well represented by exposures around Eagle Bluff Lookout (SEL and SEB; Fig. 8A). Many of the soils, quartz dunes, and eolianite units contain a mix of *B. costulatus* (Lamarck) and *B. onslowi* (Cox) (G. Kendrick, personal communication, 2005) in at least 14 distinct levels, labelled in italics “a” to “n” stratigraphically from the base to the top of the section (Fig. 8A).

Fig. 8. (A) Litho- and aminostratigraphy from central Shark Bay. Lower case italic alphabet define lithostratigraphic units from the oldest “a” to the youngest “n” in the basin. Average *Bothriembryon* A/I s from upper units are shown with calibrated AMS ^{14}C ages (from Table 3). A MIS 5e correlated Aminozone E (~125 ka) at Eagle Bluff is based on A/I correlation with nearby Baba Head (Table 4). (B) Approximate parabolic curve showing evolution of A/I in *Bothriembryon* over the past 130 ka from dated samples. The estimated ages of older units are calculated from the APK model (Mitterer and Kriausakul, 1989).

4.5. Quantification of loss of amino acids with age: increasing then decreasing A/I

Based primarily on morphostratigraphic and vertical stratigraphic succession, and limited radiometric ages, we independently ranked the relative ages of lithologic units from Shark Bay and the Zuytdorp Cliffs covering the entire depositional interval discussed in this paper. The key sections include Eagle Lookout (SEL, SEB), and Zuytdorp Cliffs (SMT and SWG) and are illustrated mostly in Figs. 8, 10 and 11. At Baba Head (SBA), MIS 5e beach deposits lap directly on to an oolite of middle Pleistocene age. Amino acid concentrations were determined based on a synthetic norleucine spike adjusted to the mass of biogenic carbonate in the samples from representative units of the lithostratigraphic succession (Table 6).

Plots of stratigraphic rank versus amino acid concentration and A/I (Table 6) quantify the progressive decrease in amino acid ($A + I$) concentration for over half of the succession as A/I increases and peaks around 0.95 and 1.20 for WR and *Bothriembryon* land snails, respectively (Fig. 9). Once the A/I attains an equilibrium value, it then decreases as

Table 6

Amino acid concentrations in whole-rock samples and *Bothriembryon* land snails from the Shark Bay/Zuytdorp Cliffs, including stratigraphic rank

Lab ID (UAL)	Sample ID	Location	A/I	Age or MIS	Concentration (nM/mg)		A + I	Straigraphic rank
					Ile	alle		
Whole rock								
4069 A	RFB 1z	Fairbridge Bluff	0.049	2010±40 yr	384.2	18.6	402.9	21
4069 B	RFB 1z	Fairbridge Bluff	0.050	1	253.4	12.6	266.0	21
4073 A	RAJ 1a	Army Jetty	0.189	9260±60 yr	31.6	6.0	37.6	20
4073 B	RAJ 1a	Army Jetty	0.165	1	77.6	12.8	90.3	20
4057 B	SEB 4l	Eagle Bluff N	0.257	2?	11.1	2.9	14.0	19
4066 A	SMT 1 (D10)	Monkey Tarra	0.417	> 19,100±130 yr	60.3	23.4	83.8	18
4121 A	SEL5k	Eagle Bluff	0.425	2–4	11.8	4.8	16.6	17
4121 B	SEL5k	Eagle Bluff	0.439	2–4	16.1	7.1	23.2	16
4068 A	SMT 4 (D10)	Monkey Tarra	0.576	> 32,250±260 yr	38.4	27.0	65.3	15
3804 A	SEB1a	Eagle Bluff	0.466	5e	28.5	13.1	41.6	14
3804 B	SEB1a	Eagle Bluff	0.509	5e	35.0	17.8	52.8	14
4129 A	SBA1d	Baba Head	0.427	122.6±0.6 ka	101.5	43.3	144.8	13
4129 B	SBA1d	Baba Head	0.595	5e	79.9	47.5	127.4	13
4059 A	SFW 6a	Fowler’s Bay Oo	0.505	5e	14.4	6.8	21.2	12
4127 A	SFW5a	Fowler’s Bay Oo	0.632	5e	5.6	3.5	9.2	12
4127 B	SFW5a	Fowler’s Bay Oo	0.565	5e	6.8	4.0	10.8	12
4058 A	SFW 2d	Fowler’s Bay Sk	0.580	5e	9.5	5.5	15.0	11
4058 B	SFW 2d	Fowler’s Bay Sk	0.693	5e	15.7	10.9	26.6	11
4130 A	SBA3b	Old Baba oolite	0.956	9 or 11	22.0	21.1	43.1	10
4130 B	SBA3b	Old Baba oolite	0.974	9 or 11	24.6	24.0	48.6	10
4130 C	SBA3b	Old Baba oolite	0.919	9 or 11	30.7	28.2	59.0	10
4130 D	SBA3b	Old Baba oolite	0.924	9 or 11	29.1	26.9	56.0	10
3813 A	PEZ1a(1)	Mt. Eliza, Perth	0	>422 ka	10.0	0.0	10.0	Not ranked
3813 B	PEZ1a(1)	Mt. Eliza, Perth	0	>422 ka	8.5	0.0	8.5	Not ranked
3814 A	PEZ2a(2)	Mt. Eliza, Perth	0	>422 ka	16.5	0.0	16.5	Not ranked
3814 B	PEZ2a(2)	Mt. Eliza, Perth	0.188	>422 ka	12.5	2.4	14.9	Not ranked
4065 A	SMT 1 (D6)	Monkey Tarra	0.745	Unknown	15.4	11.5	26.9	9
4065 B	SMT 1 (D6)	Monkey Tarra	0.859	Unknown	21.3	18.3	39.7	9
4063 A	SMT 2 (D6)	Monkey Tarra	0.670	Unknown	11.2	7.5	18.8	8
4063 B	SMT 2 (D6)	Monkey Tarra	0.754	Unknown	17.9	13.5	31.5	8
4064 A	SMT1 (D5)	Monkey Tarra	0.474	Unknown	7.3	3.5	10.8	7
4064 B	SMT1 (D5)	Monkey Tarra	0.576	Unknown	8.7	5.0	13.7	7
4062 A	SMT 2 (D4)	Monkey Tarra	0.069	Unknown	72.7	5.1	77.8	6
4062 B	SMT 2 (D4)	Monkey Tarra	0.090	Unknown	59.5	5.3	64.7	6
4116 A	SWG1 (D9b)	Woomerangee	0.456	Unknown	4.4	2.1	6.5	5
4116 B	SWG1 (D9b)	Woomerangee	0.508	Unknown	8.1	4.2	12.4	5
4115 A	SWG1 (D9a)	Woomerangee	0.152	Unknown	12.6	2.1	14.7	4
4115 B	SWG1 (D9a)	Woomerangee	0.220	Unknown	10.5	2.5	13.0	4
4117 A	SWG1 (D8)	Woomerangee	0.337	Unknown	3.9	1.3	5.2	3
4117 B	SWG1 (D8)	Woomerangee	0.421	Unknown	2.9	1.2	4.1	3
4118 B	SWG1 (D4)	Woomerangee	0.079	Unknown	24.8	2.0	26.7	2
4119 A	SWG1 (D3)	Woomerangee	0.117	Unknown	13.1	1.5	14.6	1
4119 B	SWG1 (D3)	Woomerangee	0.098	Unknown	13.9	1.4	15.3	1
Bothriembryon land snail								
4211 A	STL 3	Tetradon Loop	0.020	Modern ¹⁴ C	684.3	13.4	697.7	25
4037 D	SMT1(D10)	Monkey Tarra	0.043	1563±138 yr				24
3953 E	SBS1a	Brickie’s Q	0.224	3860±183 yr	137.0	30.7	167.7	23
3952 C	SMT4(S10)	Monkey Tarra	0.147	6370±94 yr	403.1	58.9	462.1	22
3962 D	SEB 4l	Eagle Bluff	0.289	10,528±200 yr	41.4	12.0	53.3	21
4053AB	SLL 1a	Little Lagoon	0.570	Non-finite ¹⁴ C	104.3	59.4	163.7	20
4013 C	SMM 2b	Monkey Mia	0.729	5a?	66.9	49.2	116.1	19
4013 B	SMM 2b	Monkey Mia	0.735	5a?	105.2	76.7	181.8	18
4014 B	SMM 1c	Monkey Mia	0.631	5a?	38.1	24.0	62.2	17
4014 D	SMM 1c	Monkey Mia	0.675	5a?	36.2	24.4	60.6	16
3955 A	SEB 1b	Eagle Bluff	0.812	5e	32.0	26.0	57.9	15
3955 B	SEB 1b	Eagle Bluff	0.829	5e	10.4	8.6	19.0	14
3959 D	SEB 4h (3)	Eagle Bluff	0.829	5e	38.3	31.7	70.0	13
3959 A	SEB 4h (3)	Eagle Bluff	0.988	5e	61.3	60.6	121.9	12
4011 A	SEL 1f (2)	Eagle Bluff	0.951	7–9	50.1	47.8	97.9	11
4011 B	SEL 1f (2)	Eagle Bluff	0.986	7–9	30.3	29.9	60.2	10

Table 6 (continued)

Lab ID (UAL)	Sample ID	Location	<i>A/I</i>	Age or MIS	Concentration (nM/mg)		<i>A + I</i>	Stratigraphic rank
					Ile	alle		
4010 A	SEL 1f (1)	Eagle Bluff	0.234	7–9	50.2	11.7	61.9	9
4009 D	SEL 1c	Eagle Bluff	0.907	≥11	22.8	20.7	43.5	8
4009 B	SEL 1c	Eagle Bluff	1.096	≥11	22.0	24.1	46.1	7
4213 B	SMM5	Monkey Mia	1.163	≥11	160.9	187.2	348.2	6
4213 A	SMM4	Monkey Mia	1.205	≥11	157.7	190.1	347.9	5
4140	STD 2a	Tetradon Loop	1.009	≥11	20.2	20.4	40.7	4
4208 B	STD 1a	Tetradon Loop	1.110	≥11	23.9	26.6	50.5	3
3965 C	SMT 2 (S6)	Monkey Tarra	0.391	Unknown	12.0	4.7	16.7	3
3965 D	SMT 2 (S6)	Monkey Tarra	0.421	Unknown	8.6	3.6	12.2	3
3965 B	SMT 2 (S6)	Monkey Tarra	0.732	Unknown	36.0	26.3	62.3	3
3965 A	SMT 2 (S6)	Monkey Tarra	0.894	Unknown	31.2	27.8	59.0	3
3965 E	SMT 2 (S6)	Monkey Tarra	0.932	Unknown	28.1	26.2	54.3	3
3963 E	SMT 1 (D6)	Monkey Tarra	0.655	Unknown	57.6	37.7	95.4	2
3963 B	SMT 1 (D6)	Monkey Tarra	0.673	Unknown	7.5	5.2	12.7	2
3963 C	SMT 1 (D6)	Monkey Tarra	0.867	Unknown	34.7	30.1	64.7	2
3963 D	SMT 1 (D6)	Monkey Tarra	0.967	Unknown	26.6	25.7	52.3	2
3954 E	SMT 2 (D5)	Monkey Tarra	0.503	Unknown	6.4	3.2	9.7	1
3954 A	SMT 2 (D5)	Monkey Tarra	0.583	Unknown	14.0	8.1	22.1	1
3954 D	SMT 2 (D5)	Monkey Tarra	0.760	Unknown	13.7	10.4	24.1	1
3954 G	SMT 2 (D5)	Monkey Tarra	0.772	Unknown	37.0	28.5	65.5	1
3954 C	SMT 2 (D5)	Monkey Tarra	0.779	Unknown	23.5	18.3	41.8	1
3954 B	SMT 2 (D5)	Monkey Tarra	0.875	Unknown	13.1	11.4	24.5	1
3954 F	SMT 2 (D5)	Monkey Tarra	0.962	Unknown	22.2	21.4	43.6	1

Note: Amino acid concentrations of alloisoleucine (alle) and isoleucine (Ile) are determined based on an internal synthetic norleucine spike, added during preparation based on mass of biogenic carbonate in the sample. Radiometric ages are listed in the previous tables. Some Rottneest Island/Perth sites are provided for comparison.

concentration drops below about 50–80 nM/mg, after which *A/I* generally becomes very low or unmeasurable. This progression is consistent between two unrelated sample materials and occurs somewhat earlier in WR samples (~rank 10) compared to land snails (~rank 6) from the same succession (Fig. 9), indicating an effective limit of the method of about 500 ± 100 ka. Therefore, these data support the conclusion that reversal of *A/I* trends toward lower values, progressing to non-detectable levels, is a reasonable time-dependent measure of relative age of limestone and fossils. This progression maps the diagenesis of amino acids in fossil material and independently confirms the stratigraphic and diagenetic evidence that the Zuytdorp Cliffs are the oldest Quaternary units in the Shark Bay area.

4.6. Stratigraphy, diagenesis, and constraints on the age(s) at the type locality of Tamala Limestone

As the type locality of the Tamala Limestone, the eolianite at Zuytdorp Cliffs should be representative of Quaternary eolianites across WA in terms of stratigraphy, age, and diagenetic characteristics. Two sections along the Zuytdorp Cliffs (Fig. 10) including the type locality were logged in 2002 and 2003 (Fig. 11). The first ($26.69273^\circ \times 113.6000^\circ$) is locally referred to as “Monkey Tarra” (SMT1/2); the etymology is uncertain, as no geographic names appear on topographic maps near the site. The section is about 55 m high (Fig. 11). The second is

located about 3 km north of Woomerangee Hill (SWG1/2; $26.8836^\circ \times 113.7485^\circ$), where the 135-m-high, is somewhat accessible. We collected samples from over half of the exposed units at each site, below which access was limited. However, we observed, recorded, and photographed the entire section in detail at each site (Fig. 10).

4.6.1. Stratigraphy of Monkey Tarra (SMT1/2)

The upper seven units analyzed at SMT, each generally 3–10 m thick (Fig. 11), are composed of medium to fine, buff-colored, moderately to well-sorted sand with variable kurtosis (Donald, 2003). Carbonate percentages in both eolianites and palaeosols range between 55% and 89%. The carbonate fraction is consistently composed of marine calcareous algae, benthic foraminifera, echinoderms, and molluscs, while the non-carbonate fraction is predominantly silt to fine quartz sand. The relatively low quartz content of the soils compared to those exposed in the adjacent Shark Bay area may be an indication of the distal and elevated position of the ridge and cliffs from continental eolian sources of quartz. Further, weathering of the predominantly carbonate surface appears to have been curtailed by the subsequent episodes of eolian deposition over multiple sea-level cycles.

Eolian bedforms indicate deposition in a subaerial environment. Bioturbation by vegetation and land animals is probably responsible in cases where bedding is destroyed. Marine bedforms were not observed at either Zuytdorp Cliffs site, but lower units likely to contain marine facies

Fig. 9. Amino acid concentration ($A+I$ in nM/mg) and A/I values in selected whole-rock (plates A and B) and *Bothriembryon* (plates C and D) samples compared to stratigraphic rank. The ranking is based on morphostratigraphic and vertical stratigraphic succession at several key sites in and near Shark Bay. Concentration decreases with older stratigraphic rank of samples, while A/I increases up to 0.95 to 1.10–1.20 (WR and land snails, respectively), then ultimately decreases to non-detectable levels. This progression of decreasing concentration is time-dependent and can be used under controlled parameters to infer the relative ages of deposits. Units at the Zuydorp Cliffs are represented by stratigraphic rank 9 to 1 for WR and 3 to 1 for land snails.

were not closely observed. The limestone is well cemented in all units except D/S10, which comprise recently active dunes in small <1–2 ha patches. The stratigraphically highest Pleistocene unit D/S6 could be broken with strong hammer blows, while the stratigraphically lowest units collected (D4, S4), were highly indurated, hammer-ringing hard, and difficult to break. The limestone rocks show minor alteration of the grains in the uppermost units. Vugginess and general recrystallization were observed in the rocks of the lower units. The diagenetic grade V (Fig. 5; Land et al., 1967) of the lower limestones suggests the D4 couplet is significantly older and more altered than units higher in the section (grades IV–V).

4.6.2. Amino acid results from Monkey Tarra (SMT)

WR samples from eolianites from SMT reveal a decrease in A/I in stratigraphically lower units (Fig. 11). The maximum individual WR ratios of 0.86 and 0.75 (with very low amino acid abundances) from the stratigraphically highest unit D6 suggest a *minimum* APK age of 300–400 ka, or possible correlation with MIS 9 or 11. In successively lower units, mean WR A/I values from units D5 and D4 yield only trace levels of amino acids, and mean A/I of 0.53 and 0.08, respectively.

On the upper, undulating ridge surface inland of the Zuydorp Cliffs, *Bothriembryon* from soils on subactive dunes yield a mean A/I of 0.05 ± 0.02 (3) (SMT1(D10)) and

Fig. 10. Images of Zuytdorp Cliffs: (A) View near Playford et al.'s (1975, 1976) type locality at Woomerangee Hill (SWG). Arrows identify persons for scale. (B) View near Woomerangee Hill. Note the deep karstic weathering of the surface rocks. (C) Stratigraphy at Monkey Tarra (SMT), located about 20 km north of Woomerangee along the cliffs.

0.17 ± 0.06 (3) (SMT4(S10)). The A/I values correspond well with calibrated AMS ^{14}C ages of 1563 ± 138 and 6370 ± 94 cal BP from the same individual snails. However, the carbonate sediment enclosing the land snails yields AMS ^{14}C ages of $22,262 \pm 263$ and $> 32,250$ (infinite) cal BP. Age estimates of 75 and 270 ka were determined by APK from the WR A/I values of 0.38 and 0.72, suggesting an age well beyond the range of ^{14}C dating. These results indicate that Holocene-age land snails colonized dunes composed of eroded late and middle Pleistocene sediments from the cliffs, and demonstrate that little Holocene sediment from the shelf or shore has been transported to the upper ridge surface. Further, they highlight the importance of an independent assessment of ^{14}C ages on carbonate.

4.6.3. Woomerangee Hill Section (SWG1/2)

The sedimentology of the upper SWG dune ("D") units is similar to SMT. Field surveys in the area indicate accretionary eolian bedforms up to +200 m. The coarse grain size

and preservation of biogenic grain textures suggest the dunes had a nearby source within a few kilometres. Even the uppermost limestone and paleosol units are highly indurated (Figs. 10 and 11), however, and break only with strong hammer blows. Many of the lower units are well indurated with nearly complete intergranular cementation and recrystallization. The exposed ridge surface near Woomerangee is a deeply (1–2 m) pitted, irregular karst surface (Fig. 10B). The diagenetic grade V throughout SWG is somewhat greater than the top of SMT, while a line of caves (Fig. 10A) at the boundary between couplets D/S7 and D/S8 accentuates the advanced stages of limestone weathering.

4.6.4. Amino acid results at Woomerangee Hill (SWG1/2)

WR AAE analyses of several units in the upper two-thirds of the section reveal samples with decreasing A/I values in stratigraphically lower units (Fig. 11): 0.48/0.22 in D9, 0.42 in D8, 0.08 in D4, and 0.11 in D3. The oldest units yield subsamples with low or non-detectable levels of

Fig. 11. Stratigraphic logs and mean A/I values from Woomerangee Hill and Monkey Tarra, Zuytdorp Cliffs. The stratigraphic complexity, advanced stage of diagenesis, non-detectable levels of amino acids, and comparisons with other Quaternary sequences suggest the mid-to-upper Zuytdorp Cliffs at Woomerangee are at least >500 ka, and possibly well over 1 Ma. Darker colors represent reddened paleosol horizons while the lighter colors are bioclastic eolianite (see Fig. 10).

amino acids (0–35 nM/mg). This decrease in A/I parallels that of SMT. In contrast, WR deposits within Shark Bay, only 13 km east of SMT, and 30 km north of SWG, yield higher concentrations, up to about 100 nM/mg $A+I$, and apparently reliable A/I values up to 0.97 (SBA3).

All *Bothriembryon* in SWG stratigraphic units were entirely recrystallized and more often dissolved *in situ*, yielding shell impressions. As a result, no *Bothriembryon* were analyzed from the SWG section. The presumed extinct *B. ridei* (Kendrick, 1978 and personal communication), from Edel Land north of Zuytdorp Cliffs produce A/I values as high as 1.11 (Table A1; Fig. 6), indicating a minimum APK age of >300 –400 ka. *B. costulatus* (Lamarck) from the base of Eagle Lookout in Shark Bay produced a near-equilibrium A/I value of 1.22, indicating an age of well over 400 ka. Eolianite and marine carbonates >780 ka exhibit some of the most extensive cave development in WA along the Leeuwin-Naturalist ridge in the Augusta karst area (Eberhard, 2004). Internal molds are the most common evidence of land snails in these rocks, however, several *B. naturalistarum* shells from these deposits yield exclusively non-detectable levels of amino acids (unpublished data) from recrystallized shells. Lacking alternative explanations, these results from WR and land snails suggest that the uppermost units at SWG and mid-to-lower units at SMT are most likely older

than the oldest deposits within Shark Bay and on neighboring Edel Land; that is, much older than 500 ka. Morphostratigraphic evidence further suggests that the development of the basin at Shark Bay could not progress without closure of its western margin by the earlier build up of the Zuytdorp ridge.

5. Discussion

5.1. A regional view of age sequences of eolianites in WA

Across the three regions representing much of coastal WA, the most commonly occurring eolianites range between MIS 1 (<6 –7 ka) and MIS 11–13 (<500 ka) based on AAE–APK age estimates (Fig. 6). Holocene beaches and dunes are ubiquitous along the coast, yet spatially separated by headlands, often of last interglacial age or older bedrock. Several late Pleistocene (Aminozone E) coral reefs exposed at Foul Bay (near Hamelin Bay), Rottnest Island, Leander Point, Dongara, Shark Bay (Tetradon), Cape Cuvier, and throughout Cape Range have been dated by U/Th (cited papers and Table 4), and provide independent APK age calibration for Aminozone E across a steep climate gradient. Middle Pleistocene ridges lie a few kilometers inland, where they are increasingly

buried by younger deposits. Early Pleistocene outcrops are rare along the coastline of WA. Where present, they generally display massive accumulations and extensive cave development.

5.2. Representative Quaternary limestone-eolianite sequences in WA

Rottneest-Perth—A complex of late Pleistocene (Aminozone E and C) and Holocene (Aminozone A) eolianite and marine deposits are dominant on Rottneest Island (Hearty, 2003; Hearty et al., 2007) (Fig. 4A). On the mainland along the Swan River estuary in Perth, late and middle Pleistocene rocks (Murray-Wallace and Kimber, 1989; Hearty, 2003) are exposed in several cliff sections (Minim Cove (Aminozone E); Peppermint Grove (Aminozone F?/G), Mt. Eliza at King's Park (> Aminozone H)) and in road cuttings and quarries south and north of Perth.

Cape Naturaliste to Cape Leeuwin—Important components of the Quaternary coastal stratigraphy are exposed between Cape Leeuwin and Cape Naturaliste. At Conto's Beach (Fig. 4B) south of Cape Freycinet to near C. Leeuwin, a well-developed Holocene sequence directly abuts an early Pleistocene coastal ridge, which contains the most extensive Quaternary cave system in WA (Eberhard, 2004). This site demonstrates how antecedent topography may create large gaps in the rock record. Late Pleistocene marine and eolian sediment is exposed at numerous sites including Cowaramup, Hamelin Bay (Fig. 4C), and Quarry Bay, the latter of which also contains outcrops of middle Pleistocene eolianite (Fig. 4D).

Guilderton and Nambung Pinnacles—The “Pinnacles” are striking pillar-like rock formations that occur 4–6 km inland of the WA coastline at several localities between Perth and Cervantes, and are the central natural attraction of Nambung National Park. These stone columns emerge several meters from the desert floor and are generally 1–3 m in diameter. At Nambung NP, the pinnacles, often partially buried by pure quartz sand (Fig. 4E), are composed of a variety of sediment types and internal structures including carbonate sand, quartz sand, thick (often >50 cm) and highly indurated micritic calcrete, red clayey soil, and rhizomorphs. Despite their spectacular and widespread occurrence in the area, little scientific research has been dedicated to understanding how and when they formed (see McNamara, 2002).

At Guilderton Quarry (Fig. 4F), pinnacles are progressively exhumed from within an eolianite sand quarry. The eolianite yields WR *A/I* values of 0.52, correlated with Aminozone H, estimated to be ca. 400 ka old. Considerably younger Aminozone E deposits (*A/I* = 0.32) are located 5 km westward at the coastline at Guilderton. The complexity and advanced stages of diagenesis of the pinnacles, manifest by massive calcretes and complex internal structure, combined with their inland morphostratigraphic position, are commensurate with an APK age estimate of ca. 400 ka. This age estimate contrasts with the youthful age of formation of 25 ka suggested by McNamara (2002).

We interpret the pinnacles as the preserved casts and molds of large trees in ancient forests, buried by the advance and aggradation of coastal eolianites. The casts create conduits for groundwater and precipitation of calcrete, which ultimately fill and become more highly indurated than the enclosing eolianite. When the eolianite is eroded, the more-resistant pillars are left standing in relief (Fig. 4EF). The process of tree burial and casting during have been described for last-interglacial eolianites in Bermuda and the Bahamas (Neumann and Hearty, 1996; Hearty and Kindler, 1997), and provides a model to explain pinnacle formation in south-central WA.

5.3. Global comparisons of the parallel diagenetic effects on amino acids and aging limestone

These geochemical and chronostratigraphic findings from both WR and land-snails are in general agreement with late Quaternary successions from Bermuda (Hearty et al., 1992), Bahamas (Hearty and Kaufman, 2000), and Hawaii (Hearty et al., 2000, 2005a, b), as well as with the apparent relative ice-volume/sea-level changes interpreted from deep-sea $\delta^{18}\text{O}$ records (Fig. 12; from Raymo et al., 1990). These studies support a trend of greater highstand amplitude over the past 0.5 Ma (MIS 13 to 1; identified as Interval “iv” in Fig. 12, with the probable exception of MIS 7) particularly during MIS 11, 9, and 5e. During Interval “iii” (0.5–1.0 Ma; Fig. 12), peak interglacial sea levels were comparatively lower than Interval “iv”. During this interval of lower highstands, carbonate sediment was either sparsely or not deposited across large areas of the coastal zone currently above sea level. Lacking carbonate input, this interval was characterized by the development of dense, red and clayey soils (aka “Big Red Soil” or BRS of Hearty et al., 2005a) on coastal limestone. Although we have yet to confirm BRS correlatives in WA, sections at Cape Peron, Broadhurst North, and Monkey Mia expose dense, deep red clayey paleosols of middle Pleistocene or older age in basal units, similar to the BRS.

Quaternary limestone successions and mean aminozone *A/I* values in Bermuda, Bahamas, and Hawaiian are comparable to SW and NW WA (Tables 5 and 7). Bermuda shares not only similar latitudes as Perth and Rottneest (~32°N), but also similar average climate and mean annual temperature (MAT) (ca. 18–20 °C) compared to SW WA. Mean WR aminozone values are nearly identical between these distant localities (Hearty, 2003; Table 7). Likewise, mean temperatures of ca. 24–25 °C and latitudes of 21–24°N latitude from the Bahamas and Hawaiian Islands (Kauai, Oahu, and Molokai) are similar to those in NW WA.

In these cases, AAE results from carbonate sequences show consistent reliable mean *A/I*s in stratigraphic units assigned to Aminozone H. Rocks older than the BRS generally show a reversal in *A/I* or non-detectable levels of amino acids (Table 7). We suspect these factors are closely linked to recrystallization and karst processes, which are almost always more advanced in early Pleistocene rocks

Fig. 12. Deep-sea isotope curve from Raymo et al. (1990) from which generalized ice-volume changes are interpreted. Several intervals (“i”–“iv”) of higher and lower relative high stands are recognized between the late Pliocene and the present. The majority of eolianite exposures and aminozones on the Western Australia coastal plain are correlated with the latest interval “iv”, while those from the type Tamala Limestone at the Zuytdorp Cliffs are correlated with interval “ii”.

predating the Brunhes at >0.78 Ma in these regions (Hearty et al., 1992, 1996). Thus, by comparison, geologic and AAE data from Bermuda, Bahamas, and Hawaii further support an *early Pleistocene* age of the Zuytdorp Cliffs, as suggested over three decades ago by Logan et al. (1970, 1974). MIS 25, 31, and 37 stand out as important interglacials (Fig. 12) during which many of these older units may have been deposited. We anticipate future Sr isotopic and paleomagnetic studies will corroborate these observations.

5.4. Highstand or lowstand eolianites?

This investigation has proceeded under the well-supported premise that eolianites represent supratidal shoreline facies deposited during peak interglacial highstands. A “peak highstand” includes three intervals: (1) the approach of sea level to its maximum position during transgression, (2) the interval when sea level stabilizes at or has minor shifts within its general maximum still stand position, and (3) the fall of sea level from its maximum position at the onset of glaciations. The configuration of the shelf and nearshore ramp, supply of sediments, onshore energy vectors, and magnitude of superimposed fluctuations on the highstand curve must also be considered along each coastline. Each interglacial has a maximum sea-level position above, at, or lower than present, during which eolian sediments can be delivered to each of these zones. Eolianites discovered by drilling far below present sea level may have been deposited during significantly lower, or much older peak sea levels, or as sediment was shifted down slope.

Our stratigraphic, sedimentological, and AAE findings support the basic assumption of predominantly highstand deposition of eolianites with the following arguments:

Uniformitarianism—Eolianites are being deposited during the current Holocene highstand in WA. Subtidal, intertidal, and supratidal facies can be observed in

shallowing-upward sequences formed during the current highstand. AAE independently demonstrates the age equivalence of subtidal, intertidal, and supratidal facies for Holocene and older interglacial deposits. U/Th-dated and AAE-correlated last-interglacial marine and eolian facies are rich in carbonate (>80% in most cases) and confirm a correlation with highstand events.

Diagenesis—Carbonate sediments, rich in aragonite, indurate rapidly under vadose conditions, and thus are not prone to migration. Once emplaced above sea level, the dunes are rapidly colonized by plants and animals, and quickly cement at grain contacts in the vadose environment.

Temporal limitations—At Eagle Bluff, Shark Bay, for example, there are a minimum of twelve terrigenous quartz dunes and soil units, but only two or three intercalated lenses of carbonate representing the stratigraphic record for the past ca. 400 ka (Donald, 2003). Considering that glacial periods comprise the majority (~90%) of the Quaternary Period, if carbonate dunes were constantly migrating across the continental shelf by “winds... considerably stronger during the glacial periods” (Playford, 1997, p. 794), deposition of a greater amount of carbonate sediment would be expected in the deposits.

Sedimentology—The coarse grain size in most eolianites suggests proximity to a high-energy shoreline. After tens of kilometres of migration, progressively finer-grained and smaller volumes of sands and silts would remain, inhibiting further dune propagation.

Geographic and topographic barriers—The distance from the lowstand shoreline to the present coastal ridges of WA ranges is between 50 and 100 km. Carbonate dunes generally do not migrate more than a few kilometers unless repeatedly disturbed by fire or overgrazing (Hearty et al., 2000). If this occurred, sediments would be transported as transverse or more likely parabolic dunes or barchanoid dunes. These dunes would probably not maintain a constant volume, but rather would be

Table 7

Proposed correlation of marine isotope stages (MIS), and aminozones of major units in WA Regions I and II with whole-rock aminostratigraphic successions from Bermuda, Bahamas, and the Hawaiian Islands

Amino-zone	WA Region I (this study) and Hearty (2003) in bold	WA Region II Shark Bay and Zuytdorp Cliffs (this study: bold)	Bermuda ^a site/Fm	Bahamas ^b New Providence, Eleuthera, and Exuma Islands	Hawaiian Islands ^c Oahu, Kauai (couplets), and Molokai Islands	MIS
A3	Modern beach-dunes	Modern beach-dunes	Modern beach $0.12 \pm 0.01(2)$	Modern beach-dunes $0.05 \pm 0.02(3)$	Modern beach-dunes $0.11 \pm 0.03(7)$ Couplet VIII	Mod; Late 1
A1–2	Unit VI	Unit m; eolianite		Windermeer I $0.10 \pm 0.01(2)$		Mid/ late 1
C/D	Rottneest upper eolianites Unit V	Unit k; eolianite	Southampton Fm $0.23 \pm 0.03(3)$	Green Cay; First Bay $0.29 \pm 0.03(5)$	$0.34 \pm 0.01(6)$ Couplet VIIb	5a/5c?
E2	Minim Cove Unit IV	Unit h; Eagle Bluff; Baba Head Bibra Fm	Rocky Bay Fm $0.27 \pm 0.03(13)$	Lyford Cay; Boiling Hole $0.38 \pm 0.02(12)$	$0.46 \pm 0.04(27)$ Couplet VIIa	Late 5e
E1	Fairbridge reef Unit IV	Gladstone North	Grape Bay Mb $0.40 \pm 0.02(13)$		$0.51 \pm 0.03(11)$ Couplet VIIa	Early 5e
F?/G	Peppermint Grove Unit III	Dampier Fm	Upper Town Hill Fm $0.49 \pm 0.04(11)$	Blue Hill; Glass Window $0.58 \pm 0.01(3)$	$0.65 \pm 0.01(2)$ Couplet VI/V	7?/9
H	Guilderton and Nambung Pinnacles Unit II	Unit a; Baba Head yellow oolite	Lower Town Hill Fm $0.67 \pm 0.03(6)$ 405 ± 5 ka	Hunt's Q; Goulding Cay $0.67 \pm 0.05(16)$ <550 ka	$0.81 \pm 0.08(6)$ Couplet IV	11
Big red soil	No known equivalent	BRS? Basal units Peron Pen	Government/ Wilkinson Quarries	Exuma cores: BRS^d	Wiopili Quarry, Kauai, BRS	12– 24?
K	Leeuwin- Naturaliste ridge ^e (e.g., Jewel Caves); all ND	Zuytdorp Cliffs units; low <i>A/I</i> and ND	Walsingham Fm Reverse Polarity $1.11 \pm 0.02(3)$; mostly ND	Exuma Cays ^d Reverse polarity; all ND	All ND in Couplets III, II, and I	25? to 37?

Note: All correlations are based on aminostratigraphy and morphostratigraphic succession. Mean *A/I* values of whole-rock samples from WA localities are listed in Table 5.

References:

^aHearty et al. (1992); Hearty and Vacher (1994); Hearty et al. (1999); Hearty (2002a).

^bHearty and Kaufman (2000).

^cHearty et al. (2000, 2005a, b).

^dHearty et al. (1996; unpublished).

^eEberhard (2004).

thinned and distributed over large areas. The finite sediment supply (disconnected from source) and high rates of sediment dispersal would greatly inhibit long-distance migration in any dune form. Moreover, after migration of 50–100 km from the shelf break, it is unlikely that the dunes would cease to migrate, and “rack up” to form a succession of shore-parallel or stacked ridges as exists along the present coastline.

The Zuytdorp Cliffs themselves present a barrier to sand migration. If the current highstand of the Holocene cannot effectively deposit modern sediment on top of the Zuytdorp Cliffs 50–100 m high, as indicated by our samples, then a coastline 120 m lower, and 75 km distant

from the present cliff would make transport and deposition of migrating dune sediment implausible.

Difficulties with preservation of lowstand dunes—Submerged shorelines and localized deposits, interpreted to be coastal dunes, have been described in proximity to glacial shorelines (James et al., 2004), but few such cases have been documented on mid-to-upper continental shelf areas of WA. If the sediment were unconsolidated, the carbonate dune morphology would effectively be destroyed by transgressing seas. If the lowstand dunes were indurated, vast fields of eolianites should be preserved across the continental shelf, which have not been detected.

5.5. Stratigraphic and AAE constraints on the ages of interbedded quartz deposits

Pits, lenses, and beds of fine-to-coarse quartz sand occur across coastal SW WA. In Region I, the differences are accentuated with nearly pure quartz deposited on nearly pure carbonate beds. Farther north in Regions II and III, quartz is more common in eolianites, due to aridity and the proximity of continental dunes. The ages of many of the quartz deposits are constrained by their association with the AAE relative ages of bounding carbonates.

Our study does not support the “residual soil” formation theory (e.g., Bastian, 1996) as applicable to coastal WA for the following reasons:

- (1) Primary dune morphology is preserved in most Aminozone A–H carbonate-rich deposits of WA capped by quartz. Primary morphology would be lost if tens of meters of eolianite were dissolved.
- (2) Eolianites retain their original pristine mineralogy (high percentage of aragonite) even where overlain by beds or localized pits of pure quartz. If quartz were being concentrated by dissolution of eolianites to become “residual soils” (Bastian, 1996), a pattern of increasing % quartz or decreasing % carbonate with age is expected. Instead, quartz beds are deposited on eolianites representing each of the aminozone deposits “C” through “H”. In a majority of cases (55%), the insoluble quartz fraction is 100% (Table 2), while underlying eolianites are often of similar richness in CaCO₃ (i.e., 80–99%; Table A1). Thus, if the underlying carbonates are pristine, the overlying pure quartz beds are unlikely to be genetically related to them. Furthermore, the 30–40-m-high, pure-quartz dunes observed today among the coastal carbonates (for example, at Mandurah Quarry (Fig. 3D), Ankatel, and Lancelin: WMQ1, WAN1, and WLN1a in Table 2) cannot be explained by eolianite dissolution!
- (3) Extensive diagenesis would be a precursor of dissolution, yet even after hundreds of thousands of years, there is no evidence of a dissolution front, increasing percentages of quartz in older deposits, or the progressive loss of carbonate in eolianite. Aminozones A–H contain no significant karst or cave development that would be expected if these deposits were extensively dissolved over to yield pure and widespread quartz beds.

The essential physical evidence to support residual soil formation is lacking from WA. The quartz deposits in WA are more likely explained by continental dunes that expanded onto the coastal plain, as postulated by Semeniuk and Glassford (1988). The purity of the interbedded quartz and carbonate deposits (Allayialis, 2003) implies out-of-phase deposition of each mineral facies, presumably during glacial lowstand and interglacial phases, respectively.

5.6. Is the Tamala-type locality representative of WA eolianites?

The preponderance of coastal eolianites sampled in WA is relatively young and datable by AAE, falling into Aminozones A through H at ≤ 500 ka (Fig. 6). Most coastal-cliff profiles contain active modern, subactive, or stabilized Holocene dunes. A significant number of exposures are associated with the broad last interglacial (MIS 5) highstand, elsewhere dated between 80 and 130 ka. This interval comprises two groups, Aminozones C and E, best exposed on Rottnest Island (Hearty, 2003). Along the Swan River estuary in Perth, facies exposures of older Aminozones F?/G and H deposits are relatively complete.

The considerable length of time required for the development of up to 10 couplets of limestone with well-developed exposure surfaces and paleosols, together with the advanced state of diagenesis, including karst and cavernous weathering, indicate the type Tamala Limestone at the Zuytdorp Cliffs is of much greater age than most Quaternary eolianite exposed in the coastal zone between Cape Leeuwin and Cape Range. The quantified decrease in the concentration of amino acids from stratigraphically lower beds at the Zuytdorp Cliffs similarly reflects an age significantly greater than Aminozone H (ca. 400 ka) from nearby Shark Bay. In Bermuda, Bahamas, and Hawaii, amino acid concentrations decrease to unmeasurable levels in many dated deposits older than Aminozone H (Table 7) and the BRS. We therefore conclude that the middle and upper beds at the type locality at the Zuytdorp Cliffs are of early Pleistocene age, and those lower in the section perhaps may be as old as late Pliocene. To avoid the confusion associated with the type Tamala Limestone at the Zuytdorp Cliffs, we suggest restricting the term “Tamala Limestone” to the rocks at the Zuytdorp Cliffs, and propose new informal formational names, members, and type localities for Quaternary limestone eolianites that can be applied more broadly to marine deposits of WA (Table 8), and are guided by age and AAE correlations in Tables 5, 7, and Table A1.

6. Conclusions

A regional AAE survey of 200 sites in WA from the Southern Ocean to the Timor Sea reveals a dominance of Holocene and last interglacial (*sensu lato*) sites near the coastline, with fewer middle Pleistocene exposures further inland. The degree of carbonate diagenesis in older ridges generally increases landward but reveals only moderate recrystallization and induration, and minimal cavernous weathering in most deposits defined from WR and land snails as Aminozones A to H. These aminozones most likely correlate with interglacials of MIS 1–11 or 13, based on APK age modelling and comparison with better-studied and dated Quaternary successions in Bermuda, Bahamas, and Hawaii.

Pure quartz sand deposits are often interbedded with the carbonates. These are not “residual soils” (the product of eolianite dissolution, leading to concentration of quartz).

Table 8
Proposed informal nomenclature and type localities of coastal carbonates of SW (Region I) and central Western Australia (Region II)

Division	Informal name and type locality	Subdivisions (members)	Description (facies; diagenetic grade; other features)	Other named correlatives South to North	Est age (ka)	MIS
Holocene	Army Jetty fm (Rottnest Island)	Modern beach-dunes	→ Fossiliferous marine and eolian; loose to lightly consolidated sand along most WA coastlines	Herschell mb (Playford 1990)	0	
		<i>Weak soil</i> Porpoise Bay mb (beach and dunes)	→ Mid-Holocene transgression to present sea level		10	1
Late Pleistocene	<i>Major Paleosol—quartz dune sand</i> Rottnest fm	Fish Hook Bay mb	→ Primarily eolian facies with 3–5 interbedded protosols with land invertebrates; lightly to firmly cemented, brown soils	Minim Cove (Murray-Wallace and Kimber, 1989)	75	5a–5c
		<i>Paleosol</i> Fairbridge Bluff mb	→ Reef and subtidal marine and eolian facies; red soil cap	Rottnest fm (Playford, 1990)	130	5e
Middle Pleistocene	<i>Major Paleosol—quartz dune sand</i> Guilderton fm (Quarry 5 km east of coast)	Peppermint Grove mb	→ Fossiliferous marine and eolian; firmly cemented sand; capping red soils and thick calcrete	Peppermint Grove (Murray-Wallace and Kimber, 1989)	190	7/9
		<i>Paleosols and quartz sand</i> Nambung mb	→ Eolian facies with dense and complex calcrete pillars or pinnacles; altered CaCO ₃ minerals.	Dampier fm (Logan et al., 1970; 1974)		11
		<i>Paleosols and quartz sand</i> Mt. Eliza mb	→ Eolian facies with altered CaCO ₃ minerals		500	13/15
Early Pleistocene	<i>Major Paleosol—quartz dune sand</i> Zuytdorp fm (former Tamala Limestone)	Woomerangee mb. Up to 10 cycles of eolianite— <i>paleosol couplets</i>	→ Dense, highly cemented eolianites with highly altered/recrystallized CaCO ₃ minerals; interbedded with quartz-rich paleosols; cavernous weathering at some levels		800?	> 25
					1800?	

Note: Ages are inferred or estimated based upon radiometric or amino acid epimerization data.

Neither extensive karst nor cavernous weathering, concomitant with dissolution to account for the abundance of pure quartz, is observed in these beds. Rather, we suggest that quartz sand is widely distributed across the coastal plain by eolian activity during regressive sea-level events. A minor northward shift of 10° of the Southern Ocean convergence during glacials could account for the reversal in prevailing wind from onshore (current and interglacial pattern) to offshore (glacial periods) across much of SW WA.

Advanced limestone diagenesis, including recrystallization and speleogenesis, such as occurs at the Zuytdorp Cliffs, corresponds with low amino acid concentrations, generally producing lowing or non-detectable levels of amino acids. Nearby sites in Shark Bay, experiencing comparable environmental conditions, yield consistently high *A/I* values

on WR samples and land snail specimens up to 300–600 ka. We quantify this progressive change and interpret local contrasts in *A/I* and amino acid (*A + I*) concentration as a function of the much greater age of the Zuytdorp Cliffs. The stratigraphic development of the sedimentary basin at Shark Bay was almost certainly aided by the earlier closure of its western margin facing the Indian Ocean.

The results of this regional investigation indicate the Tamala Limestone at the Zuytdorp Cliffs does not accurately represent “typical” exposed coastal deposits of WA, and therefore is undesirable as a type locality.

New informal names and type localities are proposed. Rottnest Island contains a near complete succession and a variety of shoreline facies of late Pleistocene to Holocene carbonates and soils (Hearty, 2003). The well-described,

shallowing-upward sequence exposed near the boathouse at Peppermint Grove along the Swan River in Perth (Murray-Wallace and Kimber, 1989; Hearty, 2003) is correlated with a younger middle Pleistocene highstand. A premier exposure of eolianite and unique morphological features (including “pinnacles”) at Guilderton, Nambung National Park, and several other localities were developed somewhat earlier in the middle Pleistocene (MIS 11?). We propose that the type locality at Zuytdorp Cliffs be used only in reference to the massive coastal accretions over several early Pleistocene and perhaps late Pliocene interglacial cycles at several localities in WA.

Acknowledgments

We are grateful to D. Allayialis, B. Kinna, and A. Donald for their assistance in this project and great individual efforts while working on their respective theses. The submitted manuscript was greatly improved and streamlined through the significant constructive efforts of two

anonymous reviewers and D. Kaufman. We greatly appreciate high quality of AAE data produced by the Amino Acid Laboratory at the Northern Arizona University (D. Kaufman, Director). AAE samples were analyzed and concentration data reduced by J. Bright and C. Orem. AMS ^{14}C age determinations were funded through an Australian Institute of Nuclear Science and Engineering grant (AINSE #03-056) to PJH. Support for O'Leary's Ph.D. Thesis, U/Th analyses, and a portion of the cost for AAE analyses was provided by an ARC Discovery Grant (DP0209059) to McCulloch, Hearty and Halliday. We appreciate the guidance of McCulloch and his supervision of coral dating at the MC-ICPMS facility at the Australia National University. We thank the Kanagawa Museum (administered by the Australian Academy of Sciences) for funding to PJH for logistics and analyses during 2002–2003.

Appendix A

See Table A1.

Table A1

Summary of amino acid data for all sites and stratigraphic levels from Western Australia ranked by site mean A/I of each sample material from each region

Site name	Field number	Ca CO ₃	Latitude (°S)	Longitude (°E)	A/I (site mean)	± 1	N	Aminozone	AAE age, ^{14}C or U/Th calibration
Region I WR									
<i>Aminozone A</i>									
Cervantes beach	WCV1z	99	No GPS	No GPS	0.069	0.005	2	A	
Cerv. Pinnacles	WCP1x	74	No GPS	No GPS	0.168	0.003	2	A1	
Greenhead	WGH2x		30.072	114.9619	0.185	0.000	2	A1	
Quarry Bay	QBY1a	83	34.367	115.1357	0.192	0.002	2	?	
Cerv. Pinnacles	WCP3x	83	No GPS	No GPS	0.205	0.001	2	A1	
Cowaramup	CMP1a	93	33.867	114.9795	0.209	0.007	2	A1	
Cosy Corner	CCR3a	95	34.263	115.0336	0.211	0.017	2	A1	
Hamelin Bay	HBV1e	99	34.224	115.0239	0.218	0.004	2	A1	
Hamelin Bay	WHB1c	98	34.224	115.0239	0.221	0.016	2	A1	
Leeman	WLE2x		29.926	114.9794	0.222	0.004	2	A1	
<i>Aminozone C</i>									
Mimim Cove	PMC2b(2)	59	32.025	115.7673	0.226	0.074	2	?	
Hamelin Bay	WHB1d	99	34.224	115.0239	0.227	0.003	2	C2	
Cowaramup	CMP2a	77	33.866	114.9750	0.228	0.015	2	C2	
Hamelin Bay	HBV2x	99	34.225	115.0238	0.231	0.004	2	C2	
Hamelin Bay	HBV1g	99	34.224	115.0239	0.236	0.007	2	C2	
Waneroo	WWN1a		31.653	115.7339	0.237	0.045	2	?	
Hamelin Bay	WHB1f	96	34.224	115.0239	0.239	0.004	2	C2	
Cosey Corner	CCR5d	95	34.264	115.0341	0.241	0.018	2	C2	
Hamelin Bay	HBV2a	99	34.225	115.0238	0.241	0.003	2	C2	
Cosy Corner	CCR3c1	73	34.263	115.0336	0.246	0.021	2	C2	
Coolimba	LCO1c	94	29.812	114.9753	0.247	0.006	2	?	
Hamelin Bay	WHB1a	98	34.224	115.0239	0.253	0.006	2	C1	
<i>Aminozone E</i>									
Mimim Cove	PMC1a	97	32.024	115.7652	0.258	0.006	2	E	
Peppy Grove	WPM1aa	68	No GPS	No GPS	0.268	0.108	2	?	
Foul Bay	MFU1b	98	34.249	115.0284	0.270	0.004	2	E	
Toms thicket	PTT2a(3)	88	32.654	115.6272	0.272	0.007	2	E	
Guilderton	PGU5c	91	31.353	115.4976	0.272	0.007	2	E	

U/Th; Table 4

Table A1 (continued)

Site name	Field number	Ca CO ₃	Latitude (°S)	Longitude (°E)	A/I (site mean)	± 1	N	Aminozone	AAE age, ¹⁴ C or U/Th calibration	
Cervantes	CCV1a(1)	99	30.526	115.0742	0.277	0.005	2	E	U/Th; Table 4	
Waneroo	WWN1c		31.653	115.7339	0.282	0.016	2	E2		
Cowaramup	CMP1g	96	33.867	114.9795	0.290	0.012	2	E		
Cosy Corner	CCR1c(2)	77	34.261	115.0331	0.303	0.005	2	E		
Lancelin	WLN2b?		31.051	115.3804	0.310	0.032	2	E		
Mimim Cove	PMC2b(1)	54	32.024	115.7652	0.316	0.021	2	E	U/Th; Table 4	
Cottesloe Beach	WCT1b	52	No GPS	No GPS	0.322	0.041	2	E2		
Guilderton	PGU5a	66	31.353	115.4976	0.324	0.029	2	E		
Quarry Bay	QBY1c	84	34.367	115.1357	0.325	0.019	2	E		
Leander Point	WLP1c	44	29.278	114.9132	0.328	0.000	2	E		
Point Peron	PPP1	85	32.326	115.4133	0.329	0.009	2	E	U/Th; Table 4	
Mimim Cove	PMC2d	71	32.024	115.7652	0.338	0.006	2	E		
Toms thicket	PTT2c	83	32.654	115.6272	0.340	0.001	2	E		
Cowaramup	CMP1d	93	33.867	114.9795	0.348	0.006	2	E		
Leeman	WLE1a		29.925	114.9794	0.349	0.017	2	E		
Leander Point	WLP1c		29.278	114.9132	0.362	0.010	2	E	U/Th; Table 4	
Coolimba	48.LCO1a	99	29.812	114.9753	0.364	0.022	2	E1		
Peppy Grove	WPM1b	76	No GPS	No GPS	0.392	0.005	2	?		
White Point	WWP1a		29.475	115.0236	0.395	0.002	2	E1		
Greenhead	WGH1a		30.072	114.9613	0.417	0.006	2	E		
White Point	WWP2a		29.549	115.0755	0.419	0.006	2	E1		
<i>Aminozone F[?]/G/H</i>										
Jurien	WJR1a		30.349	115.0701	0.460	0.059	2	F/G	> 350 ± 50ka 640 ± 100 ka	
Lancelin	WLN2a		31.051	115.3804	0.473	0.008	2	F/G		
Fremantle fort	FRA1	86	No GPS	No GPS	0.478	0.000	2	> F [?] /G		
Cliff head	DCH1a	66	29.516	114.9966	0.491	0.069	2	F/G		
Jurien	WJR2a		30.442	115.0854	0.499	0.000	2	F/G		
Guilderton	PGU1a	83	31.33	115.5103	0.523	0.024	2	H		
QB Quarry	QBY2a	96	34.359	115.1330	0.712	0.002	2	> H		
Peppy Grove	WPM1a	69	No GPS	No GPS	0.470	0.131	4	?		
Mt Eliza	PEZ1a	19,35	31.9631	115.8423	ND	ND	2	?		
> 422 ka (TL)										
Rottne Island (Hearty, 2003)										
<i>Aminozone A</i>										
Fairbridge Bluff	RFB1z	93	32.0160	115.5121	0.049	0.000	2	A	¹⁴ C; Table 3	
L. Astrg Bay W	RLA2a	57	31.9918	115.5053	0.097	0.007	2	A	¹⁴ C; Table 3	
Fairbridge Bluff	RFB1x	90	32.0160	115.5121	0.126	0.001	2	A		
Rottne Bore	RRB1a	99	31.9990	115.5385	0.133	0.007	2	A		
Army Jetty	RAJ	98	32.0040	115.5039	0.120	0.000	2	A	¹⁴ C; Table 3	
Herschell	RHQ1cd	90	31.9959	115.5250	0.089	0.005	4	A		
Quarry										
Herschell	RHQ1a(2)	98	31.9959	115.5250	0.131	0.005	4	A	¹⁴ C; Table 3	
Quarry										
<i>Aminozone C</i>										
Herschell dune	RHE2	97	31.9959	115.5251	0.187	0.000	2	C2		
Herschell Hill	RHE1a	98	31.9955	115.5258	0.210	0.007	2	C2		
Henrietta	RHT1a	98	31.9955	115.5258	0.221	0.006	2	C2		
Rocks										
Lake Vincent	RLV1a	99	32.0009	115.5126	0.210	0.017	2	C2		
Narrow Neck	RNN1a	48	32.0153	115.4749	0.171	0.000	2	C2		
Abraham's Bay	RAB1a	97	200m N	Of RNN	0.195	0.000	2	C2		
Oliver Hill base	ROL1a	97	32.0071	115.5158	0.229	0.002	2	C2		
Oliver Hill RR	ROL2a	99	32.0109	115.5158	0.211	0.000	2	C2		
Parakeet Bay	RPA1a	71	31.9990	115.5121	0.201	0.016	2	C2		
The Basin	RTB1a (2)	90	31.9908	115.5333	0.207	0.008	2	C2		
Armstrong Hill	RAR1a	95	31.9946	115.5039	0.206	0.004	2	C2		
South Point	RSP1e	98	32.0212	115.4688	0.196	0.008	2	C2		
South Point	RSP1c	98	32.0212	115.4688	0.218	0.002	2	C2		
Kingston Brks	RKB1a	99	32.0043	115.5558	0.171	0.004	2	C2		
Crayfish Rock	RCF1c	92	No GPS	No GPS	0.234	0.005	2	C2		
Crayfish Rock	RCF1a	99	No GPS	No GPS	0.230	0.022	2	C2		

Table A1 (continued)

Site name	Field number	Ca CO ₃	Latitude (°S)	Longitude (°E)	A/I (site mean)	± 1	N	Aminozone	AAE age, ¹⁴ C or U/Th calibration
L. Astrg Bay E	RLA1e	95	31.9924	115.5051	0.235	0.004	2	C2	¹⁴ C; Table 3
L Astrg Bay E	RLA1c	95	31.9924	115.5051	0.276	0.008	2	C1	
Fish Hook Bay	RFH1g	99	32.0246	115.4494	0.227	0.002	2	C2	
Fish Hook Bay	RFH1e	98	32.0246	115.4494	0.243	0.008	2	C1	
Fish Hook Bay	RFH1c	94	32.0246	115.4494	0.263	0.007	2	C1	
City of York	RCY1c	99	31.9969	115.4941	0.203	0.006	2	C2	
City of York	RCY1a	81	31.9969	115.4941	0.254	0.016	2	C1	
Bathurst Light	RBH2c	94	100 m W	Of RHB1	0.252	0.012	2	C1	
Bathurst Point	RBH1a	93	31.9917	115.5408	0.312	0.019	2	>C?	
Fairbridge Bluff	RFB4x	96	0.5 km E	Of RFB1	0.232	0.007	2	C2	
Fairbridge Bluff	RFB5a	98	Q at road	Near RFB1	0.222	0.012	2	C2	
Fairbridge Bluff	RFB1c	82	32.0160	115.5121	0.245	0.021	2	C1	
Jeannie's LO	RJL1c	97	32.0272	115.5224	0.261	0.080	2	C1	
Mary Cove East	RMB2g	97	200 m E	Of RMB1	0.165	0.004	4	C2	
Mary Cove East	RMB2e	95	200 m E	Of RMB1	0.200	0.008	2	C2	
Mary Cove West	RMB1e	90	32.0191	115.4909	0.280	0.004	2	C1	
Aminozone E									
Fairbridge Bluff	RFB1a	98	32.0160	115.5121	0.344	0.080	4	E	U/Th; Table 4
Jeannie's LO	RJL2a	97	0.5 km N	Of RJL1	0.371	0.008	2	E	
Mary Cove West	RMB1c	99	32.0191	115.4909	0.339	0.056	4	E	
Mary Cove West	RMB1a	96	32.0191	115.4909	0.340	0.018	2	E	
Region II WR									
Aminozone A									
Hamelin Pool	SHP1x		No GPS	No GPS	0.057	0.000	2	A	
Geraldton	WGE2x		28.799	114.6186	0.064	0.007	2	A	
Broadhurst N	SBN1c(1)	55	No GPS	No GPS	0.064	0.001	1	A	
Aminozone C									
Eagle Bluff	SEB4l	77	26.0818	113.5782	0.257	0.001	2	A	
Geraldton	WGE1x		28.789	114.6209	0.261	0.007	2	C	
Quobba	LQB1	56	25.364639	114.4069	0.264	0.023	2	C	
Cape Burney	GCB4x	64	28.869	114.6329	0.292	0.014	2	C	
Goulet Bluff	SGT5b		26.19824	113.6834	0.293	0.070	4	<C?	
Wooramel	SWR1		No GPS	No GPS	0.293	0.030	2	<C?	
Aminozone E									
Tetradon	STD1x		No GPS	No GPS	0.415	0.008	2	<E?	U/Th; Table 4
Eagle Lookout	SEL5k		26.0818	113.5782	0.432	0.010	2	<E?	
Cape Cuvier N	LCC1h	61	24.215806	113.4184	0.482	0.025	2	E	
Cape Burney	GCB1c(3)	56	28.869	114.6329	0.484	0.034	2	E	
Eagle Bluff	SEB1a	28	No GPS	No GPS	0.487	0.030	2	E	
Baba Head	SBA1d		26.60264	113.6904	0.490	0.070	4	E	
Cape Burney	GCB1d	79	28.869	114.6329	0.494	0.086	2	E	
Fowler's Bay	SFW6a	69	26.11042	113.6224	0.505	0.044	2	E	
Eagle Lookout	SEL4k		26.0818	113.5782	0.507	0.030	2	E	
Cape Burney	GCB1c(1)	68	28.869	114.6329	0.522	0.041	2	E	
Hose Q	KHO1	14	28.103583	114.2204	0.530	0.030	2	>E?	296 ± 40 ka
Gladstone N	SGN1a		No GPS	No GPS	0.591	0.020	2	E1	
Fowler's Bay	SFW5a		No GPS	No GPS	0.599	0.050	2	E	
Cape Cuvier	LCV2.5f		24.22556	113.3914	0.621	0.067	2	E	
Fowler's Bay	SFW2d	57	26.10651	113.6191	0.636	0.080	1	E	
Cape Cuvier	LCV7f		24.23446	113.3926	0.642	0.006	2	E	
Gladstone N	SGN1b		No GPS	No GPS	0.661	0.008	2	E1	
Whalebone	SWB1c(3)	54	26.13003	113.6401	0.671	0.111	1	E1	
Aminozone G and H									
Cape Burney	GCB1a	79	28.869	114.6329	0.685	0.014	2	H	460 ± 50 ka
Quobba Quarry	LQQ1	64	24.356139	113.4114	0.817	0.025	2	G/H?	
Baba Head	SBA3b		26.60468	113.6920	0.965	0.010	2	H	

Table A1 (continued)

Site name	Field number	Ca CO ₃	Latitude (°S)	Longitude (°E)	A/I (site mean)	± 1	N	Aminozone	AAE age, ¹⁴ C or U/Th calibration
<i>Monkey Tarra</i>									
Monkey Tarra	SMT1D10	84	26.69273	113.60042	0.417	0.013	2	> C?	
Monkey Tarra	SMT4D10	78	26.69481	113.60254	0.375	0.059	2	> C?	
Monkey Tarra	SMT2(D4)	74	26.69481	113.60254	0.080	0.081	2	Unk	
Monkey Tarra	SMT2(D6)	59	26.69481	113.60254	0.712	0.072	2	Unk	
Monkey Tarra	SMT1(D5)	81	26.69273	113.60042	0.802	0.006	2	Unk	
Monkey Tarra	SMT1(D5)	60	26.69273	113.60042	0.525	0.010	2	Unk	
Monkey Tarra	SMT4 D10	89	26.6786	113.60561	0.717	0.018	2	Unk	
Monkey Tarra	SMT4 D10	96	26.69481	113.60254	0.576	0.013	2	Unk	
<i>Woomerangee Hill</i>									
Woomerangee	SWG1(D9)		26.8836	113.7485	0.221	0.059	4	Unk	
Woomerangee	SWG1D9		26.8837	113.7485	0.482	0.040	2	Unk	
Woomerangee	SWG1(D8)		26.8837	113.7485	0.421	0.100	2	Unk	
Woomerangee	SWG1(D4)		26.8837	113.7485	0.079	0.004	2	Unk	
Woomerangee	SWG1(D3)		26.8837	113.7485	0.107	0.010	2	Unk	
Region III WR									
<i>Aminozone A</i>									
Coral Bay	WCB2z	71	No GPS	No GPS	0.045	0.010	2	A	
Coral Bay	WCE1a	70	No GPS	No GPS	0.066	0.001	2	A	
Port Smith	BSM1a(1)	54	18.45508	121.81384	0.097	0.020	2	A	
Port Smith	BSM1c	64	18.45508	121.81384	0.11	0.009	2	A	
Port Smith	BSM1a(2)	51	18.45508	121.81384	0.111	0.027	2	A	¹⁴ C; Table 3
Pinnacles young dunes in NP	WCP5e	38	No GPS	No GPS	0.120	0.016	2	A	
Port Smith	BSM1z	73	18.45508	121.81384	0.144	0.068	2	A	
Jacobesz	WJZ2z	91	21.82397	114.08483	0.157	0.005	2	A	
Cape Vlaming	WVL2x	79	No GPS	No GPS	0.224	0.013	2	A	
<i>Aminozone E</i>									
Jacobesz	WJZ1e	90	21.82397	114.08483	0.556	0.064	2	E	U/Th; Table 4
Mangrove Bay	WMV1a	88	No GPS	No GPS	0.661	0.077	2	E	U/Th; Table 4
Coral Bay	WCB1x	86	No GPS	No GPS	0.670	0.026	2	E	
<i>Aminozone G/H and H</i>									
Pilgrimana	NPG1a		22.19268	113.86925	0.710	0.060	2	G/H	
Pilgrimana	NPG1c		22.19268	113.86925	0.792	0.030	2	G/H	
Coral Bay	WCR1a	80	No GPS	No GPS	0.794	0.024	2	G/H	
Tanatbiddi	WTI1a	95	No GPS	No GPS	0.829	0.008	2	G/H	
Exmouth	WEX1a	70	No GPS	No GPS	0.895	0.048	2	H	≥ 325 ± 50 ka
<i>Marine shells (Aminozone as indicated)</i>									
Broadhurst Bight	SBH1a(2)	Cer	25.5437	113.4901	0.825	0.013	2	E	
Talleifer Isthmus	STF2	Cer	26.15414	113.71074	0.728	0.035	2	E	
Baba Head	SBA1d	Cer	26.60264	113.69037	0.734	0.006	2	E	U/Th; Table 4
Broadhurst Bight	SBH1a(1)	Brah	25.5437	113.4901	0.029	0.003	2	A	
Baba Head	SBA1d	Brac	26.60264	113.69037	1.003	0.034	2	E	U/Th; Table 4
Goulet Bluff	SGT5b	Brac	26.19824	113.68341	0.984	0.080	2	E	
Tetradon Loop	STD4a	Brac	No GPS	No GPS	0.868	0.006	1	E	
Monkey Mia	SMM4	Brac	No GPS	No GPS	1.180	0.030	2	F?	
Region I Bothriembryon (Aminozone as indicated)									
Blossom Beach	ABB2b		34.4572	119.3640	0.756	0.045	3	H?	
Esperance	ENI4e		33.8377	121.9187	0.634	0.063	2	G/H?	
QB Quarry	QBY2b		34.359	115.1330	0.590	0.053	3	> H?	
Cosy Corner	CCR4e		34.266	115.0341	0.407	0.052	3	E	
Foul Bay	MFU1d		34.249	115.0284	0.337	0.077	4	E	U/Th; Table 4
Hamelin Bay	WHB1e		34.224	115.0239	0.096	0.005	3	A	
Hamelin Bay	WHB1b		34.224	115.0239	0.283	0.005	3	E	
Guilderton	PGU5b		31.353	115.4976	0.591	0.049	2	E	

Table A1 (continued)

Site name	Field number	Ca CO ₃	Latitude (°S)	Longitude (°E)	A/I (site mean)	± 1	N	Aminozone	AAE age, ¹⁴ C or U/Th calibration
Herschell dune	RHE 1b		31.9959	115.5251	0.176	0.000	0	A	
City of York	RCY 1b		31.9969	115.4941	0.330	0.020	2	C	¹⁴ C; Table 3
Region II Shark Bay Bothriembryon									
<i>Aminozone A</i>									
Tamala Hwy	STL 3		No GPS	No GPS	0.021	0.001	2	A	¹⁴ C; Table 3
Monkey Tarra	SMT1D10		26.69273	113.60042	0.051	0.016	3	A	¹⁴ C; Table 3
Monkey Tarra	SMT4 S10		26.6786	113.6056	0.167	0.063	3	A	¹⁴ C; Table 3
Brickies Sd Pit	SBS1a	15	25.92952	113.5440	0.172	0.062	5	A	
<i>Aminozone B</i>									
Goulet Bluff	SGT1c		26.20547	113.6888	0.244	0.071	3	B	
Eagle Bluff	SEB4l	77	26.0818	113.5782	0.271	0.037	4	B	
Eagle Bluff	SEB4l	77	26.0818	113.5782	0.287	0.025	3	B	
<i>Aminozone C–E</i>									
Little lagoon	SLL1a	59	25.90531	113.5440	0.570	0.007	1	C–E	
Monkey Mia	SMM1c	5	25.794611	113.7190	0.655	0.061	4	C–E	
Monkey Mia	SMM2b		25.794611	113.7190	0.680	0.091	3	C–E	
Monkey Mia	SMM2b		26.794611	114.7190	0.732	0.004	2	C–E	
Eagle Bluff	SEB1d		80 m E of SEL4	80 m E of SEL4	0.732	0.023	3	C–E	
Eagle Lookout	SEL 4k		26.0818	113.5782	0.763	0.044	2	C–E	
Eagle Bluff	SEB4h(2)	14	26.0818	113.5782	0.791	0.121	4	C–E	
Eagle Lookout	SEL2h	28	26.08832	113.5780	0.791	0.083	2	C–E	
<i>Aminozone E</i>									
Eagle Lookout	SEL3h	32	26.0818	113.5782	0.813	0.116	4	E	
Eagle Bluff	SEB1b	19	80 m E of SEL4	80 m E of SEL4	0.827	0.004	2	E	
Eagle Bluff	SEB1b	19	80 m E of SEL4	80 m E of SEL4	0.840	0.064	4	E	
Eagle Bluff	SEB4h(3)	14	26.0818	113.5782	0.846	0.101	4	E	
<i>Aminozones F–H</i>									
Eagle Lookout	SEL1c	11	26.09179	113.5789	0.865	0.430	4	> F	
Tetradon	#3 STD 5		No GPS	No GPS	0.867	0.050	2	F–H	
Eagle Lookout	SEL1f(2)	30	26.09179	113.5789	0.952	0.034	3	F–H	
Tetradon	#1 STD 1a		No GPS	No GPS	0.989	0.170	2	F–H	
Tetradon	#2 STD 2a		No GPS	No GPS	1.009	0.017	2	F–H	
Eagle Lookout	SEL1c	11	26.09179	113.5789	1.073	0.156	3	> G/H	
Eagle Lookout	SEL1c	11	26.09179	113.5789	1.156	0.084	2	> G/H	
<i>Monkey Tarra</i>									
Monkey Tarra	SMT1D10)		26.69273	113.60042	0.051	0.016	3	A	
Monkey Tarra	SMT4S10)		26.6786	113.60561	0.167	0.063	3	A	
Monkey Tarra	SMT2(S6)		26.69481	113.60254	0.674	0.256	5	?	
Monkey Tarra	SMT2(S6)		26.69481	113.60254	0.678	0.049	3	?	
Monkey Tarra	SMT2(D5)		26.69481	113.60254	0.788	0.127	7	?	
Monkey Tarra	SMT1(D6)		26.69273	113.60042	0.790	0.152	5	?	
Monkey Tarra	SMT2(D6)		26.69481	113.60254	0.830	0.087	5	?	
Monkey Tarra	SMT2(D6)		26.69481	113.60254	0.853	0.101	3	?	
Monkey Tarra	SMT1(D6)		26.69273	113.60042	0.917	0.071	2	?	

Total: 212 sites, 485 analyses, 333 WR, 137 Bothrie, 15 marine shells

Note: AAE = age based on amino acid epimerization and apparent parabolic kinetic model. U/Th ages are in Table 4 and Stirling et al. (1995, 1998; references therein). ¹⁴C ages are in Table 3 and Playford (1997).

Editorial Handling by: D. Kaufman

References

- Allayialis, D., 2003. The sedimentary characteristics of peri-continental dune deposits in Western Australia. Honours Thesis, James Cook University, Townsville, QLD, 89pp.
- Bastian, L.V., 1996. Residual soil mineralogy and dune subdivision, Swan Coastal Plain, Western Australia. *Australian Journal of Earth Sciences* 43, 31–44.
- Bretz, J.H., 1960. Bermuda: a partially drowned late mature Pleistocene karst. *Geological Society of America Bulletin* 71, 1729–1754.
- Brooke, B., 2001. The distribution of carbonate eolianite. *Earth Science Reviews* 55, 135–164.
- Cheng, H., Edwards, R.L., Hoff, J., Gallup, C.D., Richards, D.A., Asmerom, Y., 2000. The half-lives of uranium-234 and thorium-230. *Chemical Geology* 169, 17–33.
- Donald, A.J., 2003. Marine and continental sediment flux in Shark Bay, Western Australia. Honours Thesis, James Cook University, Townsville, Queensland, 104pp.
- Eberhard, S., 2004. Ecology and hydrology of a threatened groundwater-dependent ecosystem: the Jewel Cave Karst System in Western Australia. Unpublished Ph.D. Thesis, Murdoch University, Perth, Western Australia, 364pp.
- Fairbridge, R.W., Teichert, C., 1953. Soil horizons and marine bands in the coastal limestones of Western Australia. *Royal Society of New South Wales Journal and Proceedings* 86, 68–87.
- Fletcher III., C.H., Murray-Wallace, C., Glenn, C.R., Sherman, C.E., Popp, B., Hessler, A., 2005. Age and origin of late Quaternary eolianite, Kaiehu Point (Moomomi), Molokai, Hawaii. *Journal of Coastal Research* 42, 97–112.
- Gill, E.D., 1968. Quaternary shorelines research in Australia and New Zealand. *Australian Journal of Science* 31 (3), 106–111.
- Goodfriend, G.A., Gould, S.J., 1996. Paleontology and chronology of two evolutionary transitions by hybridization in the Bahamian land snail *Cerion*. *Science* 274, 1894–1897.
- Goodfriend, G.A., Stipp, J.J., 1983. Limestone and the problem of radiocarbon dating of land-snail shell carbonate. *Geology* 11, 575–577.
- Hearty, P.J., 1998. The geology of Eleuthera Island, Bahamas: a Rosetta Stone of Quaternary stratigraphy and sea-level history. *Quaternary Science Reviews* 17, 333–355.
- Hearty, P.J., 2002a. Revision of the late Pleistocene stratigraphy of Bermuda. *Sedimentary Geology* 153 (1–2), 1–21.
- Hearty, P.J., 2002b. The Kaena highstand on Oahu, Hawaii: Further evidence of Antarctic Ice collapse during the middle Pleistocene. *Pacific Science* 56 (1), 65–82.
- Hearty, P.J., 2003. Stratigraphy and ages of fossil carbonate deposits of Rottneest Island, Western Australia. *Quaternary Research* 60, 211–222.
- Hearty, P.J., Kaufman, D.S., 2000. Whole-rock aminostratigraphy and Quaternary sea-level history of the Bahamas. *Quaternary Research* 54, 163–173.
- Hearty, P.J., Kindler, P., 1997. The stratigraphy and surficial geology of New Providence Island, Bahamas. *Journal of Coastal Research* 13 (3), 798–812.
- Hearty, P.J., Vacher, H.L., 1994. Quaternary stratigraphy of Bermuda: A high-resolution pre-Sangamonian rock record. *Quaternary Science Reviews* 13, 685–697.
- Hearty, P.J., Miller, G.H., Stearns, C., Szabo, B.J., 1986. Aminostratigraphy of Quaternary shorelines around the Mediterranean basin. *Geological Society of America Bulletin* 97, 850–858.
- Hearty, P.J., Vacher, H.L., Mitterer, R.M., 1992. Aminostratigraphy and ages of Pleistocene limestones of Bermuda. *Geological Society of America Bulletin* 104, 471–480.
- Hearty, P.J., McNeill, D.F., Land, L., Dill, R.F., 1996. Stratigraphy and ages of two 33 m long bank-margin cores from the Exuma Cays, Bahamas. In: *Proceedings and Abstracts, American Association of Petroleum Geologists, San Diego, CA*, A63.
- Hearty, P.J., Kindler, P., Cheng, H., Edwards, R.L., 1999. A +20 m middle Pleistocene sea-level highstand (Bermuda and the Bahamas) due to partial collapse of Antarctic ice. *Geology* 27, 375–378.
- Hearty, P.J., Kaufman, D.S., Olson, S.L., James, H.F., 2000. Stratigraphy and whole-rock amino acid geochronology of key Holocene and Last Interglacial carbonate deposits in the Hawaiian Islands. *Pacific Science* 54 (4), 423–442.
- Hearty, P.J., O'Leary, M.J., Kaufman, D.S., Page, M., Bright, J., 2004a. Amino acid geochronology of individual foraminifer (*Pulleniatina obliquiloculata*) tests, north Queensland margin, Australia: a new approach to correlating and dating Quaternary tropical marine sediment cores. *Paleoceanography* 19, 4022–4036.
- Hearty, P.J., Olson, S.L., Kaufman, D.S., Edwards, R.L., Cheng, H., 2004b. Stratigraphy and geochronology of pitfall accumulations in caves and fissures, Bermuda. *Quaternary Science Reviews* 23, 1151–1171.
- Hearty, P.J., Karner, D.B., Renne, P.R., Olson, S.L., Fletcher, S., 2005a. $^{40}\text{Ar}/^{39}\text{Ar}$ age of a young rejuvenation basalt flow: Implications for the duration of volcanism and the timing of carbonate platform development during the Quaternary on Kaua'i, Hawaiian Islands. *New Zealand Journal of Geology and Geophysics* 48, 199–211.
- Hearty, P.J., James, H., Olson, S.L., 2005b. The geological context of paleo-lakes and Pleistocene birds at Ulupau Head, O'ahu, Hawaiian Islands. In: Alcover, J.A., Bover, P. (Eds.), *Insular Vertebrate Evolution: the Palaeontological Approach. Monografies de la Societat d'Història Natural de les Balears*, September 2003, Mallorca, Spain, pp. 113–128 (390pp).
- Hearty, P.J., O'Leary, M.J., Donald, A., Lachlan, T., 2006. The enigma of 3400 cal yr BP coastal oolites in Tropical northwest Western Australia...why then, why there? *Sedimentary Geology* 186, 171–185.
- Hearty, P.J., Hollin, J.T., Neumann, A.C., O'Leary, M.J., McCulloch, M., 2007. Global sea-level fluctuations during the last interglaciation (MIS 5e). *Quaternary Science Reviews*, doi:10.1016/j.quascirev.2007.06.019.
- James, N.P., Collins, L.B., Bone, Y., Hallock, P., 1999. Subtropical carbonates in a temperate realm: modern sediments on the southwest Australian shelf. *Journal of Sedimentary Research* 69 (6), 1297–1321.
- James, N.P., Bone, Y., Kyser, T.K., Dix, G.R., Collins, L.B., 2004. The importance of changing oceanography in controlling late Quaternary carbonate sedimentation on a high-energy, tropical, oceanic ramp: north-western Australia. *Sedimentology* 51, 1179–1205.
- Kendrick, G.W., 1978. New species of fossil nonmarine molluscs from Western Australia and evidence of late Quaternary climate change in the Shark Bay district. *Journal of the Royal Society of Western Australia* 60 (2), 49–60.
- Kendrick, G.W., Wyrwoll, K.-H., Szabo, B.J., 1991. Pliocene–Pleistocene coastal events and history along the western margin of Australia. *Quaternary Science Reviews* 10, 419–439.
- Kindler, P., Hearty, P.J., 1996. Carbonate petrology as an indicator of climate and sea-level changes: new data from Bahamian Quaternary units. *Sedimentology* 43, 381–399.
- Kinna, B., 2002. Biogenic sediment flux on Rottneest Island and the Perth region. Unpublished Honours Thesis, James Cook University, Townsville, Queensland, Australia.
- Land, L.S., Mackenzie, F.T., Gould, S.J., 1967. The Pleistocene history of Bermuda. *Geological Society of America Bulletin* 78, 993–1006.
- Logan, B.W., Read, J.F., Davies, G.R., 1970. History of carbonate sedimentation, Quaternary Epoch, Shark Bay, Western Australia. In: *Carbonate Sedimentation and Environments, Shark Bay, Western Australia, American Association of Petroleum Geologists Memoir*, pp. 38–84.
- Logan, B.W., Read, J.F., Hagan, G.M., Hoffman, P., Brown, R.G., Woods, P.J., Gebelein, C.D., 1974. Evolution and diagenesis of Quaternary carbonate sequences, Shark Bay, Western Australia. *AAPG Memoir* 22, 358.
- McCulloch, M.T., Esat, T., 2000. The coral record of Last Interglacial sea levels and sea surface temperatures. *Chemical Geology* 169, 107–129.
- McNamara, K., 2002. The Pinnacles. Western Australia Museum, Perth, 25pp.

- Miller, G.H., Brigham-Grette, J., 1989. Amino acid geochronology: resolution and precision in carbonate fossils. *Quaternary International* 1, 111–128.
- Mitterer, R.M., Kriausakul, N., 1989. Calculation of amino acid racemization ages based apparent parabolic kinetics. *Quaternary Science Reviews* 8, 353–357.
- Munsell Soil Color Charts (2002 revised ed.). Macbeth Division of Kollmorgan Instruments Corporation, New Windsor, NY 12553.
- Murray-Wallace, C.V., Kimber, R.W.L., 1989. Quaternary marine aminostratigraphy: Perth Basin, Western Australia. *Australian Journal of Earth Science* 36, 553–568.
- Murray-Wallace, C.V., Brooke, B.P., Cann, J.H., Belperio, A.P., Bourman, R.P., 2001. Whole-rock aminostratigraphy of the Coorong, Coastal Plain, South Australia: towards a 1 million year record of sea-level highstands. *Journal of the Geological Society of London* 158, 111–124.
- Neumann, A.C., Hearty, P.J., 1996. Rapid sea-level changes at the close of the Last Interglacial (Stage 5e) recorded in Bahamian Island geology. *Geology* 24, 775–778.
- O'Leary, M.J., 2007. The stratigraphy and geochronology of emergent fossil reef deposits of Western Australia: Unpublished Ph.D. Thesis, James Cook University, Townsville, Australia, 170pp.
- O'Leary, M.J., Hearty, P.J., McCulloch, M.T. U-series evidence for widespread reef development in Shark Bay during the last interglacial. *Palaeogeography, Palaeoclimatology and Palaeoecology*, in press.
- O'Leary, M.J., Hearty, P.J., McCulloch, M.T. Morphostratigraphic evidence for unstable sea levels during marine isotope stage 5e: Cape Cuvier, Western Australia. *Quaternary Research*, in review.
- Playford, P.E., 1990. Geology of the Shark Bay area, Western Australia. In: Berry, P.F., Bradshaw, S.D., Wilson, B.R. (Eds.), *Research in Shark Bay*, Report of the France–Australie Bicentenary Expedition Committee. Western Australian Museum, Perth, pp. 13–31.
- Playford, P.E., 1997. Geology and hydrology of Rottnest Island, Western Australia. In: Vacher, H.L., Quinn, T. (Eds.), *Geology and Hydrology of Carbonate Islands*, pp. 783–810.
- Playford, P.E., Cope, R.N., Cockbain, A.E., Low, G.H., Lowry, D.C., 1975. Phanerozoic, in geology of Western Australia. Western Australia Geological Survey Memoire 2, 223–433.
- Playford, P.E., Cockbain, A.E., Low, G.H., 1976. Geology of the Perth Basin, Western Australia. Western Australia Geological Survey Bulletin 124, 311pp.
- Price, D.M., Brooke, B.P., Woodroffe, C.D., 2001. Thermoluminescence dating of eolianites from Lord Howe Island and south-west Western Australia. *Quaternary Science Reviews* 20, 841–846.
- Raymo, M.E., Ruddiman, W.F., Shackleton, N.J., Oppo, D.W., 1990. Evolution of Atlantic-Pacific $\delta^{13}\text{C}$ gradients over the last 2.5 m.yr. *Earth Planetary Science Letters* 97, 353–368.
- Reimer, P.J., et al., 2004. IntCal04 terrestrial radiocarbon age calibration, 26–0 ka BP. *Radiocarbon* 46, 1029–1058.
- Roof, S., 1997. Comparison of isoleucine epimerization and leaching potential in the molluscan genera *Astarte*, *Macoma*, and *Mya*. *Geochimica et Cosmochimica Acta* 61 (24), 5325–5333.
- Sayles, R.W., 1931. Bermuda during the Ice Age. *American Academy of Arts and Sciences* 66, 381–468.
- Semeniuk, V., Glassford, D.K., 1988. Significance of eolian limestone lenses in quartz sand formations: an interdigitation of coastal and continental facies, Perth Basin, southwestern Australia. *Sedimentary Geology* 57, 199–209.
- Stirling, C.H., Esat, T.M., McCulloch, M.T., Lambeck, K., 1995. High-precision U/Th; Tab 4 dating of corals from Western Australia and implications for the timing and duration of the Last Interglacial. *Earth and Planetary Science Letters* 135, 115–130.
- Stirling, C.H., Esat, T.M., Lambeck, K., McCulloch, M.T., 1998. Timing and duration of the Last Interglacial: evidence for a restricted interval of widespread coral reef growth. *Earth and Planetary Science Letters* 160, 745–762.
- Stuvier, M., Reimer, P.J., Bard, E., Beck, J.W., Burr, G.S., Hughen, K.A., Kromer, B., McCormac, G., van der Plicht, J., Spurk, M., 1998. INTCAL 98 radiocarbon age calibration, 24,000–0 cal BP. *Radiocarbon* 40, 1041–1083.
- Stuiver, M., Reimer, P.J., Reimer, R.W., 2005. CALIB 5.0.2. [www program and documentation].
- Teichert, C., 1950. Late Quaternary changes of sea-level at Rottnest Island, Western Australia. *Proceedings of the Royal Society of Victoria* 59, 63–79.
- Vacher, H.L., Hearty, P.J., 1989. History of stage-5 sea level in Bermuda with new evidence of a rise to present sea level during substage 5a. *Quaternary Science Reviews* 8, 159–168.