

MATH141 – Autumn 2008

Outline Solutions to Tutorial Sheet – Week 13

1. If $x = 0$ and $y = 0$, then $2 \times 0 - 2 \times 0 + z = 4$, so $z = 4$; hence a point on the plane is $(0, 0, 4)$.
A normal to the plane is $(2, -2, 1)$. Therefore

$$\begin{aligned} \text{Distance} &= ((1, -2, 3) - (0, 0, 4)) \cdot \frac{(2, -1, 1)}{\sqrt{4+4+1}} \\ &= \frac{5}{3} \text{ units} \end{aligned}$$

2.

$$\begin{aligned} \mathcal{I} &= \int \left(\frac{12}{x} + x^5 \right) dx, \\ &= 12 \ln x + \frac{x^6}{6} + x, \end{aligned}$$

where c is an arbitrary constant.

3. (a) Any plane containing the line of intersection of \mathcal{P}_1 and \mathcal{P}_2 will be given by

$$x + y + z - 2 + \lambda(y - z + 1) = 0.$$

Now the point $A(1, 0, 0)$ lies on \mathcal{P}_3 , so it must satisfy the equation for the plane:

$$1 + 0 + 0 - 2 + \lambda(0 - 0 + 1) = 0 \Rightarrow -1 + \lambda = 0 \Rightarrow \lambda = 1.$$

Hence, the plane \mathcal{P}_3 is given by

$$x + y + z - 2 + (y - z + 1) = 0 \Rightarrow x + 2y - 1 = 0$$

Therefore, $\mathcal{P}_3 : x + 2y - 1 = 0$.

- (b) The angle between \mathcal{P}_1 and \mathcal{P}_2 is the angle between their normals.

$$\mathcal{P}_1 \text{ has normal } \tilde{n}_1 = (1, 1, 1); \quad \mathcal{P}_2 \text{ has normal } \tilde{n}_2 = (0, 1, -1)$$

$$\text{Now, } \cos \theta = \frac{\tilde{n}_1 \cdot \tilde{n}_2}{\|\tilde{n}_1\| \|\tilde{n}_2\|} \quad \text{and} \quad \tilde{n}_1 \cdot \tilde{n}_2 = (1, 1, 1) \cdot (0, 1, -1) = 0.$$

$$\Rightarrow \theta = \frac{\pi}{2}; \quad \text{i.e., the normals are perpendicular to each other.}$$

Therefore, $\mathcal{P}_1 \perp \mathcal{P}_2$.

4.

$$\mathcal{I} = \int \frac{1}{x\sqrt{10-x^2}} dx,$$

Use formula 22 from the table of integrals

$$\int \frac{1}{x\sqrt{a^2-x^2}} dx = -\frac{1}{a} \ln \left| \frac{a + \sqrt{a^2-x^2}}{x} \right| + c$$

Here $a = \sqrt{10}$. Thus

$$\mathcal{I} = -\frac{1}{\sqrt{10}} \ln \left| \frac{\sqrt{10} + \sqrt{10-x^2}}{x} \right| + c,$$

where c is an arbitrary constant.

5. In parametric form, the equations of the line are $x = 3 + 8t$, $y = 4 + 5t$ and $z = -3 - t$.

Substituting into the equation of the plane, we have $(3 + 8t) - 3(4 + 5t) + 5(-3 - t) = 12$, giving $t = -3$.

The point of intersection is therefore $(3 - 24, 4 - 15, -3 - (-3)) = (-21, -11, 0)$.

6.

$$\mathcal{I} = \int_e^{3e} \sqrt{x^2 + e^2} dx.$$

Use formula 28 from the table of integrals.

$$\int \sqrt{x^2 \pm e^2} dx = \frac{1}{2}x\sqrt{x^2 \pm e^2} \pm e^2 \ln |x + \sqrt{x^2 \pm e^2}| + c.$$

We have $a = e$. First of all write down the indefinite integral.

$$\int \sqrt{x^2 + e^2} dx = \frac{1}{2}x\sqrt{x^2 + e^2} + e^2 \ln |x + \sqrt{x^2 + e^2}| + c.$$

Thus

$$\begin{aligned} \mathcal{I} &= \left[\frac{1}{2}x\sqrt{x^2 + e^2} + e^2 \ln |x + \sqrt{x^2 + e^2}| \right]_e^{3e}, \\ &= \left(\frac{3e}{2}\sqrt{10e^2} + \frac{1}{2}e^2 \ln |3e + \sqrt{10e^2}| \right) - \left(\frac{e}{2}\sqrt{2e^2} + \frac{1}{2}e^2 \ln |e + \sqrt{2e^2}| \right), \\ &= \left(\frac{3\sqrt{10}}{2}e^2 + \frac{1}{2}e^2 \ln |[3 + \sqrt{10}]e| \right) - \left(\frac{\sqrt{2}}{2}e^2 + \frac{1}{2}e^2 \ln |[1 + \sqrt{2}]e| \right), \\ &= \frac{1}{2} (3\sqrt{10} - \sqrt{2}) e^2 + \frac{1}{2}e^2 \ln \left(\frac{3 + \sqrt{10}}{1 + \sqrt{2}} \right). \end{aligned}$$

7.

$$\mathcal{I} = \int \frac{1}{c \ln c} dc,$$

Use the substitution $u = \ln c$. Then $\frac{du}{dc} = \frac{1}{c}$ and consequently $dc = cdu$.

$$\begin{aligned} \mathcal{I} &= \int \frac{1}{u} du, \\ &= \ln |u| + m, \\ &= \ln |\ln c| + m, \end{aligned}$$

where m is an arbitrary constant.

8.

$$\begin{aligned}\mathcal{I} &= \int \tan x \, dx \\ &= \int \frac{\sin x}{\cos x} \, dx,\end{aligned}$$

Let $u = \cos x$,

$$\text{Then } \frac{du}{dx} = -\sin x \Rightarrow dx = -\frac{du}{\sin x},$$

$$\begin{aligned}\text{Thus } \mathcal{I} &= \int \frac{\sin x}{u} \cdot -\frac{du}{\sin x}, \\ &= -\int \frac{1}{u} \, du, \\ &= -\ln |u| + c, \\ &= -\ln |\cos x| + c, \\ &= \ln |\sec x| + c,\end{aligned}$$

where c is an arbitrary constant.