

Ultragraph C^* -algebras via topological quivers

by

TAKESHI KATSURA (Yokohama), PAUL S. MUHLY (Iowa City, IA),
AIDAN SIMS (Wollongong) and MARK TOMFORDE (Houston, TX)

Abstract. Given an ultragraph in the sense of Tomforde, we construct a topological quiver in the sense of Muhly and Tomforde in such a way that the universal C^* -algebras associated to the two objects coincide. We apply results of Muhly and Tomforde for topological quiver algebras and of Katsura for topological graph C^* -algebras to study the K -theory and gauge-invariant ideal structure of ultragraph C^* -algebras.

1. Introduction. Our objective in this paper is to show how the theory of *ultragraph C^* -algebras*, first proposed by Tomforde in [13, 14], can be formulated in the context of topological graphs [6] and topological quivers [11] in a fashion that reveals the K -theory and ideal theory (for gauge-invariant ideals) of these algebras. The class of graph C^* -algebras has attracted enormous attention in recent years. The graph C^* -algebra associated to a directed graph E is generated by projections p_v associated to the vertices v of E and partial isometries s_e associated to the edges e of E . Graph C^* -algebras, which, in turn, are a generalization of the Cuntz–Krieger algebras of [2], were first studied using groupoid methods [9, 8]. An artifact of the initial groupoid approach is that the original theory was restricted to graphs which are *row-finite* and have *no sinks* in the sense that each vertex emits at least one and at most finitely many edges ⁽¹⁾.

The connection between Cuntz–Krieger algebras and graph C^* -algebras is that each directed graph can be described in terms of its edge matrix, which is a $\{0, 1\}$ -matrix indexed by the edges of the graph; a 1 in the (e, f)

2000 *Mathematics Subject Classification*: Primary 46L05.

Key words and phrases: C^* -algebras, ultragraphs, topological graphs.

The first author was supported by JSPS, the second author was supported by NSF Grant DMS-0070405, the third author was supported by the Australian Research Council, and the fourth author was supported by NSF Postdoctoral Fellowship DMS-0201960.

⁽¹⁾ A refinement of the analysis in [9], due to Paterson [12], extends the groupoid approach to cover non-row-finite graphs. One can find a groupoid approach that handles sinks and extends the whole theory to higher rank graphs in [4]. A groupoid approach to ultragraph C^* -algebras may be found in [10].

entry indicates that the range of e is equal to the source of f . The term *row-finite* refers to the fact that in any row of the edge matrix of a row-finite graph, there are at most finitely many non-zero entries.

The two points of view, graph and matrix, led to two versions of Cuntz–Krieger theory for non-row-finite objects. In [5], C^* -algebras were associated to arbitrary graphs in such a way that the construction agrees with the original definition in the row-finite case. In [3], C^* -algebras—now called Exel–Laca algebras—were associated to arbitrary $\{0, 1\}$ -matrices, once again in such a way that the definitions coincide for row-finite matrices. The fundamental difference between the two classes of algebras is that a graph C^* -algebra is generated by a collection containing a partial isometry for each edge and a projection for each vertex, while an Exel–Laca algebra is generated by a collection containing a partial isometry for each row in the matrix (and in the non-row-finite case there are rows in the matrix corresponding to an infinite collection of edges with the same source vertex). Thus, although these two constructions agree in the row-finite case, there are C^* -algebras of non-row-finite graphs that are not isomorphic to any Exel–Laca algebra, and there are Exel–Laca algebras of non-row-finite matrices that are not isomorphic to the C^* -algebra of any graph [14].

In order to bring graph C^* -algebras of non-row-finite graphs and Exel–Laca algebras together under one theory, Tomforde introduced the notion of an ultragraph and described how to associate a C^* -algebra to such an object [13, 14]. His analysis not only brought the two classes of C^* -algebras under one rubric, but also it showed that there are ultragraph C^* -algebras that belong to neither of these classes. Ultragraphs are basically directed graphs in which the range of each edge is a non-empty *set* of vertices rather than a single vertex—thus in an ultragraph each edge points from a single vertex to a set of vertices, and directed graphs are the special case where the range of each edge is a singleton set. Many of the fundamental results for graph C^* -algebras, such as the well-known Cuntz–Krieger Uniqueness Theorem and the Gauge-Invariant Uniqueness Theorem, can be proven in the setting of ultragraphs [13]. However, other results, such as K -theory computations and ideal structure, are less obviously amenable to traditional graph C^* -algebra techniques.

Recently, Katsura [6] and Muhly and Tomforde [11] studied the notions of topological graphs and topological quivers, respectively. These structures consist of second countable locally compact Hausdorff spaces E^0 and E^1 of vertices and edges respectively with range and source maps $r, s : E^1 \rightarrow E^0$ which satisfy appropriate topological hypotheses. The main point of difference between the two (apart from a difference in edge-direction conventions) is that in a topological graph the source map is assumed to be a local hom-

eomorphism so that $s^{-1}(v)$ is discrete, whereas in a topological quiver the range map (remember the edge-reversal!) is only assumed to be continuous and open, and a system $\lambda = \{\lambda_v\}_{v \in E^0}$ of Radon measures λ_v on $r^{-1}(v)$ satisfying some natural conditions (see [11, Definition 3.1]) is supplied as part of the data. It is worth pointing out that given E^0 , E^1 , r and s , with r open, such a system of Radon measures will always exist. A topological graph can be regarded as a topological quiver by reversing the edges and taking each λ_v to be counting measure; the topological graph C^* -algebra and the topological quiver C^* -algebra then coincide. One can regard an ordinary directed graph as either a topological graph or a topological quiver by endowing the edge and vertex sets with the discrete topology, and then the topological graph C^* -algebra and topological quiver algebra coincide with the original graph C^* -algebra.

In this article we show how to build a topological quiver $\mathcal{Q}(\mathcal{G})$ from an ultragraph \mathcal{G} in such a way that the ultragraph C^* -algebra $C^*(\mathcal{G})$ and the topological quiver algebra $C^*(\mathcal{Q}(\mathcal{G}))$ coincide. We then use results of [6] and [11] to compute the K -theory of $C^*(\mathcal{G})$, to produce a listing of its gauge-invariant ideals, and to provide a version of condition (K) under which all ideals of $C^*(\mathcal{G})$ are gauge-invariant.

It should be stressed that the range map in $\mathcal{Q}(\mathcal{G})$ is always a local homeomorphism, so $\mathcal{Q}(\mathcal{G})$ can equally be regarded as a topological graph; indeed, our analysis in some instances requires results regarding topological graphs from [6] that have not yet been generalized to topological quivers. We use the notation and conventions of topological quivers because the edge-direction convention for quivers in [11] is compatible with the edge-direction convention for ultragraphs [13, 14].

The paper is organized as follows: In Section 2 we describe the commutative C^* -algebra $\mathfrak{A}_{\mathcal{G}} \subset C^*(\mathcal{G})$ generated by the projections $\{p_A : A \in \mathcal{G}^0\}$. In Section 3 we provide two alternative formulations of the defining relations among the generators of an ultragraph C^* -algebra which will prove more natural in our later analysis. In Section 4 we describe the spectrum of $\mathfrak{A}_{\mathcal{G}}$. We use this description in Section 5 to define the quiver $\mathcal{Q}(\mathcal{G})$, show that its C^* -algebra is isomorphic to $C^*(\mathcal{G})$, and compute its K -theory in terms of the structure of \mathcal{G} using results from [6]. In Section 6 we use the results of [11] to produce a listing of the gauge-invariant ideals of $C^*(\mathcal{G})$ in terms of the structure of \mathcal{G} , and in Section 7 we use a theorem of [7] to provide a condition on \mathcal{G} under which all ideals of $C^*(\mathcal{G})$ are gauge-invariant.

2. The commutative C^* -algebra $\mathfrak{A}_{\mathcal{G}}$ and its representations. For a set X , let $\mathcal{P}(X)$ denote the collection of all subsets of X .

DEFINITION 2.1. An *ultragraph* $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$ consists of a countable set of vertices G^0 , a countable set of edges \mathcal{G}^1 , and functions $s: \mathcal{G}^1 \rightarrow G^0$ and $r: \mathcal{G}^1 \rightarrow \mathcal{P}(G^0) \setminus \{\emptyset\}$.

DEFINITION 2.2. For an ultragraph $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$, we denote by $\mathfrak{A}_{\mathcal{G}}$ the C^* -subalgebra of $\ell^\infty(G^0)$ generated by the point masses $\{\delta_v : v \in G^0\}$ and the characteristic functions $\{\chi_{r(e)} : e \in \mathcal{G}^1\}$.

Let us fix an ultragraph $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$, and consider the representations of $\mathfrak{A}_{\mathcal{G}}$.

DEFINITION 2.3. For a set X , a subcollection \mathcal{C} of $\mathcal{P}(X)$ is called a *lattice* if

- (i) $\emptyset \in \mathcal{C}$,
- (ii) $A \cap B \in \mathcal{C}$ and $A \cup B \in \mathcal{C}$ for all $A, B \in \mathcal{C}$.

An *algebra* is a lattice \mathcal{C} that also satisfies the additional condition

- (iii) $A \setminus B \in \mathcal{C}$ for all $A, B \in \mathcal{C}$.

DEFINITION 2.4. For an ultragraph $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$, we let \mathcal{G}^0 denote the smallest algebra in $\mathcal{P}(G^0)$ containing the singleton sets and the sets $\{r(e) : e \in \mathcal{G}^1\}$.

REMARK 2.5. In [13], \mathcal{G}^0 was defined to be the smallest lattice—not algebra—containing the singleton sets and the sets $\{r(e) : e \in \mathcal{G}^1\}$. The change to the above definition causes no problem when defining Cuntz–Krieger \mathcal{G} -families (see the final paragraph of Section 3). Furthermore, this new definition is convenient for us in a variety of situations: It relates \mathcal{G}^0 to the C^* -algebra $\mathfrak{A}_{\mathcal{G}}$ described in Proposition 2.6, it allows us too see immediately that the set $r(\lambda, \mu)$ of Definition 2.8 is in \mathcal{G}^0 , and—most importantly—it aids in our description of the gauge-invariant ideals in Definition 6.1 and Lemma 6.2. For additional justification for the change in definition, we refer the reader to [10, Section 2].

PROPOSITION 2.6. We have $\mathcal{G}^0 = \{A \subset G^0 : \chi_A \in \mathfrak{A}_{\mathcal{G}}\}$ and

$$(2.1) \quad \mathfrak{A}_{\mathcal{G}} = \overline{\text{span}}\{\chi_A : A \in \mathcal{G}^0\}.$$

Proof. We begin by proving (2.1). Since $\{A \subset G^0 : \chi_A \in \mathfrak{A}_{\mathcal{G}}\}$ is an algebra containing $\{v\}$ and $r(e)$, we have $\mathcal{G}^0 \subset \{A \subset G^0 : \chi_A \in \mathfrak{A}_{\mathcal{G}}\}$. Hence

$$\mathfrak{A}_{\mathcal{G}} \supset \overline{\text{span}}\{\chi_A : A \in \mathcal{G}^0\}.$$

Since \mathcal{G}^0 is closed under intersections, the set $\overline{\text{span}}\{\chi_A : A \in \mathcal{G}^0\}$ is closed under multiplication, and hence is a C^* -algebra containing $\{\delta_v\}$ and $\{\chi_{r(e)}\}$. Hence $\mathfrak{A}_{\mathcal{G}} \subset \overline{\text{span}}\{\chi_A : A \in \mathcal{G}^0\}$, establishing (2.1).

We must now show that $\mathcal{G}^0 = \{A \subset G^0 : \chi_A \in \mathfrak{A}_{\mathcal{G}}\}$. We have already seen $\mathcal{G}^0 \subset \{A \subset G^0 : \chi_A \in \mathfrak{A}_{\mathcal{G}}\}$. Let $A \subset G^0$ with $\chi_A \in \mathfrak{A}_{\mathcal{G}}$. Then by (2.1),

$\|\chi_A - \sum_{k=1}^m z_k \chi_{A_k}\| < 1/2$ for some $A_1, \dots, A_m \in \mathcal{G}^0$ and $z_1, \dots, z_m \in \mathbb{C}$; moreover, since \mathcal{G}^0 is an algebra, we may assume that $j \neq k$ implies that $A_j \cap A_k = \emptyset$. But then $x \in A$ if and only if $x \in A_k$ for some (unique) k with $|1 - z_k| < 1/2$. That is, $A = \bigcup_{|1 - z_k| < 1/2} A_k \in \mathcal{G}^0$. ■

DEFINITION 2.7. A *representation* of a lattice \mathcal{C} on a C^* -algebra \mathfrak{B} is a collection of projections $\{p_A\}_{A \in \mathcal{C}}$ in \mathfrak{B} satisfying $p_\emptyset = 0$, $p_A p_B = p_{A \cap B}$, and $p_{A \cup B} = p_A + p_B - p_{A \cap B}$ for all $A, B \in \mathcal{C}$.

When \mathcal{C} is an algebra, the last condition of a representation can be replaced by the equivalent condition that $p_{A \cup B} = p_A + p_B$ for all $A, B \in \mathcal{C}$ with $A \cap B = \emptyset$.

Note that we define representations of lattices here, rather than just of algebras, so that our definition of $C^*(\mathcal{G})$ agrees with the original definition given in [13] (see the final paragraph of Section 3).

DEFINITION 2.8. For a fixed ultragraph $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$ we define

$$X = \{(\lambda, \mu) : \lambda, \mu \text{ are finite subsets of } \mathcal{G}^1 \text{ with } \lambda \cap \mu = \emptyset \text{ and } \lambda \neq \emptyset\}.$$

For $(\lambda, \mu) \in X$, we define $r(\lambda, \mu) \subset G^0$ by

$$r(\lambda, \mu) := \bigcap_{e \in \lambda} r(e) \setminus \bigcup_{f \in \mu} r(f).$$

DEFINITION 2.9. Let $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$ be an ultragraph. A collection of projections $\{p_v\}_{v \in G^0}$ and $\{q_e\}_{e \in \mathcal{G}^1}$ is said to satisfy *condition (EL)* if the following hold:

- (1) the elements of $\{p_v\}_{v \in G^0}$ are pairwise orthogonal,
- (2) the elements of $\{q_e\}_{e \in \mathcal{G}^1}$ pairwise commute,
- (3) $p_v q_e = p_v$ if $v \in r(e)$, and $p_v q_e = 0$ if $v \notin r(e)$,
- (4) $\prod_{e \in \lambda} q_e \prod_{f \in \mu} (1 - q_f) = \sum_{v \in r(\lambda, \mu)} p_v$ for all $(\lambda, \mu) \in X$ such that $|r(\lambda, \mu)| < \infty$.

From a representation of \mathcal{G}^0 , we get a collection satisfying condition (EL). We prove a slightly stronger statement.

LEMMA 2.10. *Let \mathcal{C} be a lattice in $\mathcal{P}(G^0)$ which contains the singleton sets and the sets $\{r(e) : e \in \mathcal{G}^1\}$, and let $\{p_A\}_{A \in \mathcal{C}}$ be a representation of \mathcal{C} . Then the collection $\{p_{\{v\}}\}_{v \in G^0}$ and $\{p_{r(e)}\}_{e \in \mathcal{G}^1}$ satisfies condition (EL).*

Proof. From the condition $p_\emptyset = 0$ and $p_A p_B = p_{A \cap B}$, it is easy to show that the collection satisfies conditions (1)–(3) in Definition 2.9. To see condition (4) let $(\lambda, \mu) \in X$ with $|r(\lambda, \mu)| < \infty$. Define $A, B \subset G^0$ by $A = \bigcap_{e \in \lambda} r(e)$ and $B = \bigcup_{f \in \mu} r(f)$. Then $A, B \in \mathcal{C}$, and from the definition of a representation, we obtain

$$p_A = \prod_{e \in \lambda} p_{r(e)}, \quad 1 - p_B = \prod_{f \in \mu} (1 - p_{r(f)}).$$

Since $r(\lambda, \mu)$ is a finite set, $r(\lambda, \mu) \in \mathcal{C}$ and $p_{r(\lambda, \mu)} = \sum_{v \in r(\lambda, \mu)} p_{\{v\}}$. Also, because $r(\lambda, \mu) = A \setminus B$, we obtain $r(\lambda, \mu) \cup B = A \cup B$ and $r(\lambda, \mu) \cap B = \emptyset$. Hence $p_{A \cup B} = p_{r(\lambda, \mu)} + p_B$. Since $p_{A \cup B} = p_A + p_B - p_{A \cap B}$, we get $p_{r(\lambda, \mu)} = p_A - p_{A \cap B}$. Therefore,

$$\sum_{v \in r(\lambda, \mu)} p_{\{v\}} = p_A - p_{A \cap B} = p_A(1 - p_B) = \prod_{e \in \lambda} p_{r(e)} \prod_{f \in \mu} (1 - p_{r(f)}). \quad \blacksquare$$

We will prove that from a collection satisfying condition (EL), we can construct a $*$ -homomorphism from $\mathfrak{A}_{\mathcal{G}}$ onto the C^* -subalgebra generated by that collection. To this end, we fix a listing $\mathcal{G}^1 = \{e_i\}_{i=1}^{\infty}$, and for each positive integer n define a C^* -subalgebra $\mathfrak{A}_{\mathcal{G}}^{(n)}$ of $\mathfrak{A}_{\mathcal{G}}$ to be the C^* -algebra generated by $\{\delta_v : v \in G^0\}$ and $\{\chi_{r(e_i)} : i = 1, \dots, n\}$. Note that the union of the increasing family $\{\mathfrak{A}_{\mathcal{G}}^{(n)}\}_{n=1}^{\infty}$ is dense in $\mathfrak{A}_{\mathcal{G}}$.

DEFINITION 2.11. Let n be a positive integer. Let $0^n := (0, \dots, 0) \in \{0, 1\}^n$. For $\omega = (\omega_1, \dots, \omega_n) \in \{0, 1\}^n \setminus \{0^n\}$, we set

$$r(\omega) := \bigcap_{\omega_i=1} r(e_i) \setminus \bigcup_{\omega_j=0} r(e_j).$$

LEMMA 2.12. Let n be a positive integer. For each $\omega \in \{0, 1\}^n$, we define $\lambda_{\omega}, \mu_{\omega} \subset \mathcal{G}^1$ by $\lambda_{\omega} = \{e_i : \omega_i = 1\}$ and $\mu_{\omega} = \{e_i : \omega_i = 0\}$. Then the map

$$\omega \mapsto (\lambda_{\omega}, \mu_{\omega})$$

is a bijection between $\{0, 1\}^n \setminus \{0^n\}$ and $\{(\lambda, \mu) \in X : \lambda \cup \mu = \{e_1, \dots, e_n\}\}$, and we have $r(\omega) = r(\lambda_{\omega}, \mu_{\omega})$.

Proof. The map $\omega \mapsto (\lambda_{\omega}, \mu_{\omega})$ is a bijection because $(\lambda, \mu) \mapsto \chi_{\lambda}$ provides an inverse, and $r(\omega) = r(\lambda_{\omega}, \mu_{\omega})$ by definition. \blacksquare

DEFINITION 2.13. We define $\Delta_n := \{\omega \in \{0, 1\}^n \setminus \{0^n\} : |r(\omega)| = \infty\}$.

LEMMA 2.14. For each $i = 1, \dots, n$, the set $r(e_i)$ is the disjoint union of the infinite sets $\{r(\omega)\}_{\omega \in \Delta_n, \omega_i=1}$ and the finite set $\bigcup_{\omega \notin \Delta_n, \omega_i=1} r(\omega)$.

Proof. First note that $r(\omega) \cap r(\omega') = \emptyset$ for distinct $\omega, \omega' \in \{0, 1\}^n \setminus \{0^n\}$. For $v \in r(e_i)$, define $\omega^v \in \{0, 1\}^n$ by $\omega_j^v = \chi_{r(e_j)}(v)$ for $1 \leq j \leq n$. Since $v \in r(e_i)$, we have $\omega^v \neq 0^n$, and $v \in r(\omega^v)$ by definition. Hence

$$r(e_i) = \bigcup_{v \in r(e_i)} r(\omega^v) = \bigcup_{\omega_i=1} r(\omega).$$

Since $r(\omega)$ is a finite set for $\omega \in \{0, 1\}^n \setminus \{0^n\}$ with $\omega \notin \Delta_n$, the result follows. \blacksquare

PROPOSITION 2.15. The C^* -algebra $\mathfrak{A}_{\mathcal{G}}^{(n)}$ is generated by $\{\delta_v : v \in G^0\} \cup \{\chi_{r(\omega)} : \omega \in \Delta_n\}$.

Proof. For each $\omega \in \Delta_n$, we have

$$\chi_{r(\omega)} = \prod_{\omega_i=1} \chi_{r(e_i)} \prod_{\omega_j=0} (1 - \chi_{r(e_j)}) \in \mathfrak{A}_{\mathcal{G}}^{(n)},$$

giving inclusion in one direction. It follows from Lemma 2.14 that the generators of $\mathfrak{A}_{\mathcal{G}}^{(n)}$ all belong to the C^* -algebra generated by $\{\delta_v : v \in G^0\} \cup \{\chi_{r(\omega)} : \omega \in \Delta_n\}$, establishing the reverse inclusion. ■

For each $\omega \in \Delta_n$, the C^* -subalgebra of $\mathfrak{A}_{\mathcal{G}}^{(n)}$ generated by $\{\delta_v : v \in r(\omega)\}$ and $\chi_{r(\omega)}$ is isomorphic to the unitization of $c_0(r(\omega))$. Since the C^* -algebra $\mathfrak{A}_{\mathcal{G}}^{(n)}$ is a direct sum of such C^* -subalgebras indexed by the set Δ_n and the C^* -subalgebra $c_0(G^0 \setminus \bigcup_{\omega \in \Delta_n} r(\omega))$ (recall that the $r(\omega)$'s are pairwise disjoint), we have the following:

LEMMA 2.16. *For two families $\{p_v\}_{v \in G^0}$ and $\{q_\omega\}_{\omega \in \Delta_n}$ of mutually orthogonal projections in a C^* -algebra \mathfrak{B} satisfying*

$$p_v q_\omega = \begin{cases} p_v & \text{if } v \in r(\omega), \\ 0 & \text{if } v \notin r(\omega), \end{cases}$$

there exists a $$ -homomorphism $\pi_n : \mathfrak{A}_{\mathcal{G}}^{(n)} \rightarrow \mathfrak{B}$ with $\pi_n(\delta_v) = p_v$ for $v \in G^0$ and $\pi_n(\chi_{r(\omega)}) = q_\omega$ for $\omega \in \Delta_n$.*

PROPOSITION 2.17. *Let $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$ be an ultragraph, and \mathfrak{B} be a C^* -algebra. Then there exist natural bijections between the following sets:*

- (i) *the set of $*$ -homomorphisms from $\mathfrak{A}_{\mathcal{G}}$ to \mathfrak{B} ,*
- (ii) *the set of representations of \mathcal{G}^0 on \mathfrak{B} ,*
- (iii) *the set of collections of projections in \mathfrak{B} satisfying condition (EL).*

Specifically, if $\pi : \mathfrak{A}_{\mathcal{G}} \rightarrow \mathfrak{B}$ is a $$ -homomorphism, then $p_A := \pi(\chi_A)$ for $A \in \mathcal{G}^0$ gives a representation of \mathcal{G}^0 ; if $\{p_A\}_{A \in \mathcal{G}^0}$ is a representation of \mathcal{G}^0 on \mathfrak{B} , then $\{p_{\{v\}}\}_{v \in G^0} \cup \{p_{r(e)}\}_{e \in \mathcal{G}^1}$ satisfies condition (EL); and if a collection of projections $\{p_v\}_{v \in G^0} \cup \{q_e\}_{e \in \mathcal{G}^1}$ satisfies condition (EL), then there exists a unique $*$ -homomorphism $\pi : \mathfrak{A}_{\mathcal{G}} \rightarrow \mathfrak{B}$ such that $\pi(\delta_v) = p_v$ and $\pi(\chi_{r(e)}) = q_e$ for all $v \in G^0$ and $e \in \mathcal{G}^1$.*

Proof. Clearly, we have the map from (i) to (ii), and by Lemma 2.10 we have the map from (ii) to (iii). Suppose that $\{p_v\}_{v \in G^0}$ and $\{q_e\}_{e \in \mathcal{G}^1}$ is a collection of projections satisfying condition (EL). Fix a positive integer n . For each $\omega \in \{0, 1\}^n \setminus \{0^n\}$, we define $q_\omega = \prod_{\omega_i=1} q_{e_i} \prod_{\omega_j=0} (1 - q_{e_j}) \in \mathfrak{B}$. Then $\{q_\omega : \omega \in \{0, 1\}^n \setminus \{0^n\}\}$ is mutually orthogonal. By Definition 2.9(3), we have

$$p_v q_\omega = \begin{cases} p_v & \text{if } v \in r(\omega), \\ 0 & \text{if } v \notin r(\omega). \end{cases}$$

Hence by Lemma 2.16, there exists a $*$ -homomorphism $\pi_n: \mathfrak{A}_{\mathcal{G}}^{(n)} \rightarrow \mathfrak{B}$ such that $\pi_n(\delta_v) = p_v$ for $v \in G^0$ and $\pi_n(\chi_{r(\omega)}) = q_\omega$ for $\omega \in \Delta_n$. For $\omega \in \{0, 1\}^n \setminus \{0^n\}$ with $|r(\omega)| < \infty$, we have $\pi_n(\chi_{r(\omega)}) = \sum_{v \in r(\omega)} p_v = q_\omega$ by Definition 2.9(4). Hence we obtain

$$\pi_n(\chi_{r(e_i)}) = \pi_n\left(\sum_{\substack{\omega \in \{0,1\}^n \\ \omega_i=1}} \chi_{r(\omega)}\right) = \sum_{\substack{\omega \in \{0,1\}^n \\ \omega_i=1}} q_\omega = q_{e_i}$$

for $i = 1, \dots, n$. Thus for each n , the $*$ -homomorphism $\pi_n: \mathfrak{A}_{\mathcal{G}}^{(n)} \rightarrow \mathfrak{B}$ satisfies $\pi_n(\delta_v) = p_v$ for $v \in G^0$ and $\pi_n(\chi_{r(e_i)}) = q_{e_i}$ for $i = 1, \dots, n$. Since there is at most one $*$ -homomorphism of $\mathfrak{A}_{\mathcal{G}}^{(n)} \rightarrow \mathfrak{B}$ with this property, the restriction of the $*$ -homomorphism $\pi_{n+1}: \mathfrak{A}_{\mathcal{G}}^{(n+1)} \rightarrow \mathfrak{B}$ to $\mathfrak{A}_{\mathcal{G}}^{(n)}$ coincides with π_n . Hence there is a $*$ -homomorphism $\pi: \mathfrak{A}_{\mathcal{G}} \rightarrow \mathfrak{B}$ such that $\pi(\delta_v) = p_v$ for $v \in G^0$ and $\pi(\chi_{r(e)}) = q_e$ for $e \in \mathcal{G}^1$. The $*$ -homomorphism π is unique because $\mathfrak{A}_{\mathcal{G}}$ is generated by $\{\delta_v : v \in G^0\} \cup \{\chi_{r(e)} : e \in \mathcal{G}^1\}$. ■

COROLLARY 2.18. *Let $\mathcal{G}^1 = \{e_i\}_{i=1}^\infty$ be some listing of the edges of \mathcal{G} . To check that a family of projections $\{p_v\}_{v \in G^0} \cup \{q_e\}_{e \in \mathcal{G}^1}$ satisfies condition (EL), it suffices to verify that conditions (1)–(3) of Definition 2.9 hold and that (4) holds for $(\lambda, \mu) \in X$ with $|r(\lambda, \mu)| < \infty$ and $\lambda \cup \mu = \{e_1, \dots, e_n\}$ for some n .*

We conclude this section by computing the K -groups of the C^* -algebra $\mathfrak{A}_{\mathcal{G}}$.

DEFINITION 2.19. For an ultragraph $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$, we denote by $Z_{\mathcal{G}}$ the (algebraic) subalgebra of $\ell^\infty(G^0, \mathbb{Z})$ generated by $\{\delta_v : v \in G^0\} \cup \{\chi_{r(e)} : e \in \mathcal{G}^1\}$.

An argument similar to the proof of Proposition 2.6 shows that

$$Z_{\mathcal{G}} = \left\{ \sum_{k=1}^n z_k \chi_{A_k} : n \in \mathbb{N}, z_k \in \mathbb{Z}, A_k \in \mathcal{G}^0 \right\}.$$

PROPOSITION 2.20. *We have $K_0(\mathfrak{A}_{\mathcal{G}}) \cong Z_{\mathcal{G}}$ and $K_1(\mathfrak{A}_{\mathcal{G}}) = 0$.*

Proof. For each $n \in \mathbb{N} \setminus \{0\}$, let $Z_{\mathcal{G}}^{(n)}$ be the subalgebra of $\ell^\infty(G^0, \mathbb{Z})$ generated by $\{\delta_v : v \in G^0\} \cup \{\chi_{r(e_i)} : i = 1, \dots, n\}$. By an argument similar to the paragraph following Proposition 2.15, we see that $Z_{\mathcal{G}}^{(n)}$ is a direct sum of the unitizations (as algebras) of $c_0(r(\omega), \mathbb{Z})$'s and $c_0(G^0 \setminus \bigcup_{\omega \in \Delta_n} r(\omega), \mathbb{Z})$. Hence the description of $\mathfrak{A}_{\mathcal{G}}^{(n)}$ in the above-mentioned paragraph shows that there exists an isomorphism $K_0(\mathfrak{A}_{\mathcal{G}}^{(n)}) \rightarrow Z_{\mathcal{G}}^{(n)}$ which sends $[\delta_v], [\chi_{r(\omega)}] \in K_0(\mathfrak{A}_{\mathcal{G}}^{(n)})$ to $\delta_v, \chi_{r(\omega)} \in Z_{\mathcal{G}}^{(n)}$. By taking inductive limits, we get an isomorphism $K_0(\mathfrak{A}_{\mathcal{G}}) \rightarrow Z_{\mathcal{G}}$ which sends $[\chi_A] \in K_0(\mathfrak{A}_{\mathcal{G}})$ to $\chi_A \in Z_{\mathcal{G}}$ for $A \in \mathcal{G}^0$.

That $K_1(\mathfrak{A}_{\mathcal{G}}) = 0$ follows from the fact that $K_1(\mathfrak{A}_{\mathcal{G}}^{(n)}) = 0$ for each n , and by taking direct limits. ■

REMARK 2.21. It is not difficult to see that the isomorphism in Proposition 2.20 preserves the natural order and scaling.

3. C^* -algebras of ultragraphs

DEFINITION 3.1. Let $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$ be an ultragraph. A vertex $v \in G^0$ is said to be *regular* if $0 < |s^{-1}(v)| < \infty$. The set of all regular vertices is denoted by $G_{\text{rg}}^0 \subset G^0$.

DEFINITION 3.2. For an ultragraph $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$, a *Cuntz–Krieger \mathcal{G} -family* is a representation $\{p_A\}_{A \in \mathcal{G}^0}$ of \mathcal{G}^0 in a C^* -algebra \mathfrak{B} and a collection of partial isometries $\{s_e\}_{e \in \mathcal{G}^1}$ in \mathfrak{B} with mutually orthogonal ranges that satisfy

- (1) $s_e^* s_e = p_{r(e)}$ for all $e \in \mathcal{G}^1$,
- (2) $s_e s_e^* \leq p_{s(e)}$ for all $e \in \mathcal{G}^1$,
- (3) $p_v = \sum_{s(e)=v} s_e s_e^*$ for all $v \in G_{\text{rg}}^0$,

where we write p_v in place of $p_{\{v\}}$ for $v \in G^0$.

The C^* -algebra $C^*(\mathcal{G})$ is the C^* -algebra generated by a universal Cuntz–Krieger \mathcal{G} -family $\{p_A, s_e\}$.

We will show that this definition of $C^*(\mathcal{G})$ and the following natural generalization of the definition of Exel–Laca algebras in [3] are both equivalent to the original definition of $C^*(\mathcal{G})$ in [13, Definition 2.7].

DEFINITION 3.3. For an ultragraph $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$, an *Exel–Laca \mathcal{G} -family* is a collection of projections $\{p_v\}_{v \in G^0}$ and partial isometries $\{s_e\}_{e \in \mathcal{G}^1}$ with mutually orthogonal ranges for which

- (1) the collection $\{p_v\}_{v \in G^0} \cup \{s_e^* s_e\}_{e \in \mathcal{G}^1}$ satisfies condition (EL),
- (2) $s_e s_e^* \leq p_{s(e)}$ for all $e \in \mathcal{G}^1$,
- (3) $p_v = \sum_{s(e)=v} s_e s_e^*$ for $v \in G_{\text{rg}}^0$.

PROPOSITION 3.4. *For each Cuntz–Krieger \mathcal{G} -family $\{p_A, s_e\}$, the collection $\{p_v, s_e\}$ is an Exel–Laca \mathcal{G} -family. Conversely, for each Exel–Laca \mathcal{G} -family $\{p_v, s_e\}$, there exists a unique representation $\{p_A\}$ of \mathcal{G}^0 on the C^* -algebra generated by $\{p_v, s_e\}$ such that $p_{\{v\}} = p_v$ for $v \in G^0$ and $\{p_A, s_e\}$ is a Cuntz–Krieger \mathcal{G} -family.*

Proof. This follows from Proposition 2.17. ■

COROLLARY 3.5. *Let $\{p_v, s_e\}$ be the Exel–Laca \mathcal{G} -family in $C^*(\mathcal{G})$. For an Exel–Laca \mathcal{G} -family $\{P_v, S_e\}$ on a C^* -algebra \mathfrak{B} , there exists a $*$ -homomorphism $\phi: C^*(\mathcal{G}) \rightarrow \mathfrak{B}$ such that $\phi(p_v) = P_v$ and $\phi(s_e) = S_e$. The $*$ -homomorphism ϕ is injective if $P_v \neq 0$ for all $v \in G^0$ and there exists a strongly*

continuous action β of \mathbb{T} on \mathfrak{B} such that $\beta_z(P_v) = P_v$ and $\beta_z(S_e) = zS_e$ for $v \in G^0$, $e \in \mathcal{G}^1$, and $z \in \mathbb{T}$.

Proof. The first part follows from Proposition 3.4, and the rest from [13, Theorem 6.8] because $\phi(p_A) \neq 0$ for all non-empty A if $\phi(p_v) = P_v \neq 0$ for all $v \in G^0$. ■

It is easy to see that Proposition 3.4 is still true if we replace \mathcal{G}^0 by any lattice contained in \mathcal{G}^0 and containing $\{v\}$ and $r(e)$ for all $v \in G^0$ and $e \in \mathcal{G}^1$ (see Lemma 2.10). Hence the restriction gives a natural bijection from Cuntz–Krieger \mathcal{G} -families in the sense of Definition 3.2 to the Cuntz–Krieger \mathcal{G} -families of [13, Definition 2.7]. Thus the C^* -algebra $C^*(\mathcal{G})$ is naturally isomorphic to the C^* -algebra defined in [13, Theorem 2.11].

4. The spectrum of the commutative C^* -algebra $\mathfrak{A}_{\mathcal{G}}$ Let $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$ be an ultragraph. In this section, we describe the spectrum of the commutative C^* -algebra $\mathfrak{A}_{\mathcal{G}}$ concretely. Fix a listing $\mathcal{G}^1 = \{e_i\}_{i=1}^{\infty}$ of the edges of \mathcal{G} . As described in the paragraph following the proof of Lemma 2.10, the C^* -algebra $\mathfrak{A}_{\mathcal{G}}$ is equal to the inductive limit of the increasing family $\{\mathfrak{A}_{\mathcal{G}}^{(n)}\}_{n=1}^{\infty}$, where $\mathfrak{A}_{\mathcal{G}}^{(n)}$ is the C^* -subalgebra of $\mathfrak{A}_{\mathcal{G}}$ generated by $\{\delta_v : v \in G^0\} \cup \{\chi_{r(e_i)} : i = 1, \dots, n\}$. In order to compute the spectrum of $\mathfrak{A}_{\mathcal{G}}$, we first compute the spectrum of $\mathfrak{A}_{\mathcal{G}}^{(n)}$ for a positive integer n .

DEFINITION 4.1. For $n \in \mathbb{N} \setminus \{0\}$, we define a topological space $\Omega_{\mathcal{G}}^{(n)}$ such that $\Omega_{\mathcal{G}}^{(n)} = G^0 \sqcup \Delta_n$ as a set and $A \sqcup Y$ is open in $\Omega_{\mathcal{G}}^{(n)}$ for $A \subset G^0$ and $Y \subset \Delta_n$ if and only if $|r(\omega) \setminus A| < \infty$ for all $\omega \in Y$.

For each $v \in G^0$, $\{v\}$ is open in $\Omega_{\mathcal{G}}^{(n)}$, and a fundamental system of neighborhoods of $\omega \in \Delta_n \subset \Omega_{\mathcal{G}}^{(n)}$ is

$$\{A \sqcup \{\omega\} : A \subset G^0, |r(\omega) \setminus A| < \infty\}.$$

Hence G^0 is a discrete dense subset of $\Omega_{\mathcal{G}}^{(n)}$. Note that $\Omega_{\mathcal{G}}^{(n)}$ is a disjoint union of the finitely many compact open subsets $r(\omega) \sqcup \{\omega\}$ for $\omega \in \Delta_n$ and the discrete set $G^0 \setminus \bigcup_{\omega \in \Delta_n} r(\omega)$. This fact and the paragraph following Proposition 2.15 show the following:

LEMMA 4.2. *There exists an isomorphism $\pi^{(n)} : \mathfrak{A}_{\mathcal{G}}^{(n)} \rightarrow C_0(\Omega_{\mathcal{G}}^{(n)})$ such that $\pi^{(n)}(\delta_v) = \delta_v$ and $\pi^{(n)}(\chi_{r(\omega)}) = \chi_{r(\omega) \sqcup \{\omega\}}$ for $v \in G^0$ and $\omega \in \Delta_n$.*

LEMMA 4.3. *For $i = 1, \dots, n$, the closure $\overline{r(e_i)}$ of $r(e_i) \subset \Omega_{\mathcal{G}}^{(n)}$ is the compact open set $r(e_i) \sqcup \{\omega \in \Delta_n : \omega_i = 1\}$, and we have $\pi^{(n)}(\chi_{r(e_i)}) = \chi_{\overline{r(e_i)}}$.*

Proof. This follows from Lemma 2.14. ■

Let $\tilde{\Delta}_n := \Delta_n \cup \{0^n\}$. We can define a topology on $\tilde{\Omega}_{\mathcal{G}}^{(n)} := G^0 \sqcup \tilde{\Delta}_n$ similarly to Definition 4.1 so that $\tilde{\Omega}_{\mathcal{G}}^{(n)}$ is the one-point compactification of $\Omega_{\mathcal{G}}^{(n)}$. The restriction map $\{0, 1\}^{n+1} \rightarrow \{0, 1\}^n$ induces a map $\tilde{\Delta}_{n+1} \rightarrow \tilde{\Delta}_n$, and hence a map $\tilde{\Omega}_{\mathcal{G}}^{(n+1)} \rightarrow \tilde{\Omega}_{\mathcal{G}}^{(n)}$. It is routine to check that this map is a continuous surjection, and the induced $*$ -homomorphism $C_0(\Omega_{\mathcal{G}}^{(n)}) \rightarrow C_0(\Omega_{\mathcal{G}}^{(n+1)})$ coincides with the inclusion $\mathfrak{A}_{\mathcal{G}}^{(n)} \hookrightarrow \mathfrak{A}_{\mathcal{G}}^{(n+1)}$ via the isomorphisms in Lemma 4.2.

For each element

$$\omega = (\omega_1, \omega_2, \dots, \omega_i, \dots) \in \{0, 1\}^\infty,$$

we define $\omega|_n \in \{0, 1\}^n$ by $\omega|_n = (\omega_1, \omega_2, \dots, \omega_n)$. The space $\{0, 1\}^\infty$ is a compact space with the product topology, and it is homeomorphic to $\varprojlim \{0, 1\}^n$.

DEFINITION 4.4. We define

$$\tilde{\Delta}_\infty := \{\omega \in \{0, 1\}^\infty : \omega|_n \in \tilde{\Delta}_n \text{ for all } n\},$$

and $\Delta_\infty := \tilde{\Delta}_\infty \setminus \{0^\infty\}$ where $0^\infty := (0, 0, \dots) \in \{0, 1\}^\infty$.

Since $\tilde{\Delta}_\infty$ is a closed subset of $\{0, 1\}^\infty$, the space Δ_∞ is locally compact, and its one-point compactification is homeomorphic to $\tilde{\Delta}_\infty$. By definition, $\tilde{\Delta}_\infty \cong \varprojlim \tilde{\Delta}_n$.

DEFINITION 4.5. We define a topological space $\Omega_{\mathcal{G}}$ as follows: $\Omega_{\mathcal{G}} = G^0 \sqcup \Delta_\infty$ as a set, and $A \sqcup Y$ is open in $\Omega_{\mathcal{G}}$ for $A \subset G^0$ and $Y \subset \Delta_\infty$ if and only if for each $\omega \in Y$ there exists an integer n satisfying

- (i) if $\omega' \in \Delta_\infty$ and $\omega'|_n = \omega|_n$, then $\omega' \in Y$,
- (ii) $|r(\omega|_n) \setminus A| < \infty$.

Equivalently, $A \sqcup Y \subset \Omega_{\mathcal{G}}$ is closed if and only if $Y \subset \Delta_\infty$ is closed in the product topology on $\{0, 1\}^\infty$, and for each $\omega \in \Delta_\infty$ with $|r(\omega|_n) \cap A| = \infty$ for all n , we have $\omega \in Y$.

We can define a topology on $\tilde{\Omega}_{\mathcal{G}} := G^0 \sqcup \tilde{\Delta}_\infty$ similarly to the definition above, so that $\tilde{\Omega}_{\mathcal{G}}$ is the one-point compactification of $\Omega_{\mathcal{G}}$ and $\tilde{\Omega}_{\mathcal{G}} \cong \varprojlim \tilde{\Omega}_{\mathcal{G}}^{(n)}$.

LEMMA 4.6. *In the space $\Omega_{\mathcal{G}}$, the closure $\overline{r(e_i)}$ of $r(e_i) \subset \Omega_{\mathcal{G}}$ is the compact open set $r(e_i) \sqcup \{\omega \in \Delta_\infty : \omega_i = 1\}$.*

Proof. This follows from the homeomorphism $\tilde{\Omega}_{\mathcal{G}} \cong \varprojlim \tilde{\Omega}_{\mathcal{G}}^{(n)}$ combined with Lemma 4.3. ■

PROPOSITION 4.7. *There exists an isomorphism $\pi: \mathfrak{A}_{\mathcal{G}} \rightarrow C_0(\Omega_{\mathcal{G}})$ such that $\pi(\delta_v) = \delta_v$ and $\pi(\chi_{r(e)}) = \chi_{\overline{r(e)}}$ for $v \in G^0$ and $e \in \mathcal{G}^1$.*

Proof. Taking the inductive limit of the isomorphisms $\pi^{(n)}$ in Lemma 4.2 produces an isomorphism

$$\pi : \mathfrak{A}_{\mathcal{G}} \rightarrow \varinjlim C_0(\Omega_{\mathcal{G}}^{(n)}) \cong C_0(\Omega_{\mathcal{G}}).$$

This isomorphism satisfies the desired condition by Lemma 4.6. ■

By the isomorphism π in the proposition above, we can identify the spectrum of $\mathfrak{A}_{\mathcal{G}}$ with the space $\Omega_{\mathcal{G}}$.

5. Topological quivers and K -groups In this section we will construct a topological quiver $\mathcal{Q}(\mathcal{G})$ from \mathcal{G} , and show that the C^* -algebra $C^*(\mathcal{G})$ is isomorphic to the C^* -algebra $C^*(\mathcal{Q}(\mathcal{G}))$ of [11]. Fix an ultragraph $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$, and define

$$\mathcal{Q}(\mathcal{G}) := (E(\mathcal{G})^0, E(\mathcal{G})^1, r_{\mathcal{Q}}, s_{\mathcal{Q}}, \lambda_{\mathcal{Q}})$$

as follows. Let $E(\mathcal{G})^0 := \Omega_{\mathcal{G}}$ and

$$E(\mathcal{G})^1 := \{(e, x) \in \mathcal{G}^1 \times \Omega_{\mathcal{G}} : x \in \overline{r(e)}\},$$

where \mathcal{G}^1 is considered as a discrete set, and $\overline{r(e)} \subset \Omega_{\mathcal{G}}$ are compact open sets (see Lemma 4.6).

We define a local homeomorphism $r_{\mathcal{Q}} : E(\mathcal{G})^1 \rightarrow E(\mathcal{G})^0$ by $r_{\mathcal{Q}}(e, x) := x$, and a continuous map $s_{\mathcal{Q}} : E(\mathcal{G})^1 \rightarrow E(\mathcal{G})^0$ by $s_{\mathcal{Q}}(e, x) := s(e) \in G^0 \subset E(\mathcal{G})^0$. Since $r_{\mathcal{Q}}$ is a local homeomorphism, $r_{\mathcal{Q}}^{-1}(x)$ is discrete and countable for each $x \in E(\mathcal{G})^0$. For each $x \in E(\mathcal{G})^0$ we define the measure λ_x on $r_{\mathcal{Q}}^{-1}(x)$ to be counting measure, and set $\lambda_{\mathcal{Q}} = \{\lambda_x : x \in E(\mathcal{G})^0\}$.

Reversing the roles of the range and source maps, we can also regard $\mathcal{Q}(\mathcal{G})$ as a topological graph $E(\mathcal{G})$ in the sense of [6], and its C^* -algebra $\mathcal{O}(E(\mathcal{G}))$ is naturally isomorphic to $C^*(\mathcal{Q}(\mathcal{G}))$ (see [11, Example 3.19]). Since some of the results about $C^*(\mathcal{Q}(\mathcal{G}))$ which we wish to apply have only been proved in the setting of [6] to date, we will frequently reference these results; the reversal of edge-direction involved in regarding $\mathcal{Q}(\mathcal{G})$ as a topological graph is implicit in these statements. We have opted to use the notation and conventions of [11] throughout, and to reference the results of [11] where possible only because the edge-direction conventions there agree with those for ultragraphs [13].

We let $E(\mathcal{G})_{\text{rg}}^0$ denote the largest open subset of $E(\mathcal{G})^0$ such that the restriction of $s_{\mathcal{Q}}$ to $s_{\mathcal{Q}}^{-1}(E(\mathcal{G})_{\text{rg}}^0)$ is surjective and proper.

LEMMA 5.1. *We have $E(\mathcal{G})_{\text{rg}}^0 = G_{\text{rg}}^0$.*

Proof. Since the image of $s_{\mathcal{Q}}$ is contained in $G^0 \subset E(\mathcal{G})^0$, we have $E(\mathcal{G})_{\text{rg}}^0 \subset G^0$. For each $v \in G^0$, we see that $v \in E(\mathcal{G})_{\text{rg}}^0$ if and only if $s_{\mathcal{Q}}^{-1}(v)$ is non-empty and compact because $\{v\}$ is open in $E(\mathcal{G})^0$. Since

$\{e\} \times \overline{r(e)} \subset E(\mathcal{G})^1$ is compact for all $e \in \mathcal{G}^1$, it follows that $s_{\mathcal{Q}}^{-1}(v) = \{(e, x) \in E(\mathcal{G})^1 : s(e) = v\}$ is non-empty and compact if and only if $\{e \in \mathcal{G}^1 : s(e) = v\}$ is non-empty and finite; that is, if and only if v is in G_{rg}^0 . ■

For the statement of the following theorem, we identify $C_0(E(\mathcal{G})^0)$ with $\mathfrak{A}_{\mathcal{G}}$ via the isomorphism in Proposition 4.7, and let $\chi_e \in C_c(E(\mathcal{G})^1)$ denote the characteristic function of the compact open subset $\{e\} \times r(e) \subset E(\mathcal{G})^1$ for each $e \in \mathcal{G}^1$. We denote by $(\psi_{\mathcal{Q}(\mathcal{G})}, \pi_{\mathcal{Q}(\mathcal{G})})$ the universal generating $\mathcal{Q}(\mathcal{G})$ -pair, and by $\{p_A^{\mathcal{G}}, s_e^{\mathcal{G}} : A \in \mathcal{G}^0, e \in \mathcal{G}^1\}$ the universal generating Cuntz–Krieger \mathcal{G} -family.

THEOREM 5.2. *There is an isomorphism from $C^*(\mathcal{G})$ to $C^*(\mathcal{Q}(\mathcal{G}))$ which is canonical in the sense that it takes $p_A^{\mathcal{G}}$ to $\pi_{\mathcal{Q}(\mathcal{G})}(\chi_A)$ and $s_e^{\mathcal{G}}$ to $\psi_{\mathcal{Q}(\mathcal{G})}(\chi_e)$ for all $A \in \mathcal{G}^0$ and $e \in \mathcal{G}^1$. Moreover, this isomorphism is equivariant for the gauge actions on $C^*(\mathcal{G})$ and $C^*(\mathcal{Q}(\mathcal{G}))$.*

Proof. It is easy to check using Lemma 5.1 and Proposition 3.4 that:

- (1) for each Cuntz–Krieger \mathcal{G} -family $\{p_A, s_e : A \in \mathcal{G}^0, e \in \mathcal{G}^1\}$ there is a unique $\mathcal{Q}(\mathcal{G})$ -pair $(\pi_{q,t}, \psi_{p,s})$ satisfying $\pi_{p,s}(\chi_A) = p_A$ for each $A \in \mathcal{G}^0$ and $\psi_{p,s}(\chi_e) = s_e$ for each $e \in \mathcal{G}^1$,
- (2) for each $\mathcal{Q}(\mathcal{G})$ -pair (π, ψ) , the formulae $p_A^{\pi, \psi} := \pi(\chi_A)$ and $s_e^{\pi, \psi} := \psi(\chi_e)$ determine a Cuntz–Krieger \mathcal{G} -family $\{p_A^{\pi, \psi}, s_e^{\pi, \psi} : A \in \mathcal{G}^0, e \in \mathcal{G}^1\}$.

The result then follows from the universal properties of the two C^* -algebras $C^*(\mathcal{G})$ and $C^*(\mathcal{Q}(\mathcal{G}))$. ■

REMARK 5.3. To prove Theorem 5.2, one could alternatively use the gauge-invariant uniqueness theorems for ultragraphs [13, Theorem 6.8] or topological graphs [6, Theorem 4.5], or topological quivers [11, Theorem 6.16].

THEOREM 5.4. *Let $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$ be an ultragraph. Let $\partial : \mathbb{Z}^{G_{\text{rg}}^0} \rightarrow Z_{\mathcal{G}}$ be defined by $\partial(\delta_v) = \delta_v - \sum_{e \in s^{-1}(v)} \chi_{r(e)}$ for $v \in G_{\text{rg}}^0$. Then $K_0(C^*(\mathcal{G})) \cong \text{coker}(\partial)$ and $K_1(C^*(\mathcal{G})) \cong \ker(\partial)$.*

Proof. Since $C_0(E(\mathcal{G})_{\text{rg}}^0) \cong c_0(G_{\text{rg}}^0)$ and $C_0(E(\mathcal{G})^0) \cong \mathfrak{A}_{\mathcal{G}}$, we have $K_0(C_0(E(\mathcal{G})_{\text{rg}}^0)) \cong \mathbb{Z}^{G_{\text{rg}}^0}$, $K_0(C_0(E(\mathcal{G})^0)) \cong Z_{\mathcal{G}}$ and $K_1(C_0(E(\mathcal{G})_{\text{rg}}^0)) = K_1(C_0(E(\mathcal{G})^0)) = 0$ by Proposition 2.20. It is straightforward to see that the map $[\pi_r] : K_0(C_0(E(\mathcal{G})_{\text{rg}}^0)) \rightarrow K_0(C_0(E(\mathcal{G})^0))$ in [6, Corollary 6.10] satisfies $[\pi_r](\delta_v) = \sum_{e \in s^{-1}(v)} \chi_{r(e)}$ for $v \in G_{\text{rg}}^0$. Hence the conclusion follows from [6, Corollary 6.10]. ■

6. Gauge-invariant ideals In this section we characterize the gauge-invariant ideals of $C^*(\mathcal{G})$ for an ultragraph \mathcal{G} in terms of combinatorial data associated to \mathcal{G} .

For the details of the following, see [11]. Let $\mathcal{Q} = (E^0, E^1, r, s, \lambda)$ be a topological quiver. We say that a subset $U \subset E^0$ is *hereditary* if, whenever $e \in E^1$ satisfies $s(e) \in U$, we have $r(e) \in U$. We say that U is *saturated* if, whenever $v \in E_{\text{rg}}^0$ satisfies $r(s^{-1}(v)) \subset U$, we have $v \in U$.

Suppose that $U \subset E^0$ is open and hereditary. Then

$$\mathcal{Q}_U := (E_U^0, E_U^1, r|_{E_U^1}, s|_{E_U^1}, \lambda|_{E_U^0})$$

is a topological quiver, where $E_U^0 = E^0 \setminus U$ and $E_U^1 = E^1 \setminus r^{-1}(U)$.

We say that a pair (U, V) of subsets of E^0 is *admissible* if

- (1) U is a saturated hereditary open subset of E^0 ,
- (2) V is an open subset of E_U^0 with $E_{\text{rg}}^0 \setminus U \subset V \subset (E_U^0)_{\text{rg}}$.

It follows from [11, Theorem 8.22] that the gauge-invariant ideals of $C^*(\mathcal{Q})$ are in bijective correspondence with the admissible pairs (U, V) of \mathcal{Q} .

Let $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$ be an ultragraph. We define admissible pairs of \mathcal{G} in a similar way to the above, and show that these are in bijective correspondence with the gauge-invariant ideals of $C^*(\mathcal{G})$.

DEFINITION 6.1. A subcollection $\mathcal{H} \subset \mathcal{G}^0$ is said to be an *ideal* if it satisfies

- (1) $A, B \in \mathcal{H}$ implies $A \cup B \in \mathcal{H}$,
- (2) $A \in \mathcal{H}$, $B \in \mathcal{G}^0$ and $B \subset A$ imply $B \in \mathcal{H}$.

Let $\pi: \mathfrak{A}_{\mathcal{G}} \rightarrow C_0(\Omega_{\mathcal{G}})$ be the isomorphism in Proposition 4.7. For an ideal \mathcal{H} of \mathcal{G}^0 , the set $\overline{\text{span}}\{\chi_A : A \in \mathcal{H}\}$ is an ideal of the C^* -algebra $\mathfrak{A}_{\mathcal{G}}$. Hence there exists an open subset $U_{\mathcal{H}}$ of $\Omega_{\mathcal{G}}$ with

$$C_0(U_{\mathcal{H}}) = \pi(\overline{\text{span}}\{\chi_A : A \in \mathcal{H}\}).$$

LEMMA 6.2. *The correspondence $\mathcal{H} \mapsto U_{\mathcal{H}}$ is a bijection from the set of all ideals of \mathcal{G}^0 to the set of all open subsets of $\Omega_{\mathcal{G}}$.*

Proof. Since $\mathfrak{A}_{\mathcal{G}}$ is an AF-algebra, every ideal of $\mathfrak{A}_{\mathcal{G}}$ is generated by its projections. From this fact, we see that $\mathcal{H} \mapsto \overline{\text{span}}\{\chi_A : A \in \mathcal{H}\}$ is a bijection from the set of all ideals of \mathcal{G}^0 to the set of all ideals of $\mathfrak{A}_{\mathcal{G}}$. Hence the conclusion follows from the well-known fact that $U \mapsto C_0(U)$ is a bijection from the set of all open subsets of $\Omega_{\mathcal{G}}$ to the set of all ideals of $C_0(\Omega_{\mathcal{G}})$. ■

REMARK 6.3. The existence of this bijection is one of the advantages of changing the definition of \mathcal{G}^0 from that given in [13].

LEMMA 6.4. *Let \mathcal{H} be an ideal of \mathcal{G}^0 , and let $U_{\mathcal{H}} \subset \Omega_{\mathcal{G}}$ be the corresponding open set. Then for $v \in G^0$, $\{v\} \in \mathcal{H}$ if and only if $v \in U_{\mathcal{H}}$, and for $e \in \mathcal{G}^1$, $r(e) \in \mathcal{H}$ if and only if $\overline{r(e)} \subset U_{\mathcal{H}}$.*

Proof. This follows from Proposition 4.7. ■

DEFINITION 6.5. We say that an ideal $\mathcal{H} \subset \mathcal{G}^0$ is *hereditary* if, whenever $e \in \mathcal{G}^1$ satisfies $\{s(e)\} \in \mathcal{H}$, we have $r(e) \in \mathcal{H}$, and that it is *saturated* if, whenever $v \in G_{\text{rg}}^0$ satisfies $r(e) \in \mathcal{H}$ for all $e \in s^{-1}(v)$, we have $\{v\} \in \mathcal{H}$.

PROPOSITION 6.6. *An ideal \mathcal{H} of \mathcal{G}^0 is hereditary (resp. saturated) if and only if the corresponding open subset $U_{\mathcal{H}} \subset \Omega_{\mathcal{G}} = E(\mathcal{G})^0$ is hereditary (resp. saturated) in the topological quiver $\mathcal{Q}(\mathcal{G})$.*

Proof. An open subset $U \subset \Omega_{\mathcal{G}} = E(\mathcal{G})^0$ is hereditary if and only if, whenever $(e, x) \in E(\mathcal{G})^1$ satisfies $s_{\mathcal{Q}}(e, x) = s(e) \in U$, we have $r_{\mathcal{Q}}(e, x) = x \in U$. This is equivalent to the statement that, whenever $e \in \mathcal{G}^1$ satisfies $s(e) \in U$, we have $\overline{r(e)} \subset U$. Thus Lemma 6.4 shows that an ideal \mathcal{H} is hereditary if and only if $U_{\mathcal{H}}$ is hereditary.

An open subset $U \subset \Omega_{\mathcal{G}} = E(\mathcal{G})^0$ is saturated if and only if, whenever $v \in E(\mathcal{G})_{\text{rg}}^0 = G_{\text{rg}}^0$ satisfies $r_{\mathcal{Q}}(s_{\mathcal{Q}}^{-1}(v)) \subset U$, we have $v \in U$. For $v \in G_{\text{rg}}^0$, we have $r_{\mathcal{Q}}(s_{\mathcal{Q}}^{-1}(v)) = \bigcup_{e \in s^{-1}(v)} \overline{r(e)}$. Hence U is saturated if and only if, whenever $v \in G_{\text{rg}}^0$ satisfies $\overline{r(e)} \subset U$ for all $e \in s^{-1}(v)$, we have $v \in U$. Thus Lemma 6.4 again shows that an ideal \mathcal{H} is saturated if and only if $U_{\mathcal{H}}$ is saturated. ■

DEFINITION 6.7. Let \mathcal{H} be a hereditary ideal of \mathcal{G}^0 . For $v \in G^0$, we define $s_{\mathcal{G}/\mathcal{H}}^{-1}(v) \subset \mathcal{G}^1$ by

$$s_{\mathcal{G}/\mathcal{H}}^{-1}(v) := \{e \in \mathcal{G}^1 : s(e) = v \text{ and } r(e) \notin \mathcal{H}\}.$$

We define $(G_{\mathcal{H}}^0)_{\text{rg}} \subset G^0$ by

$$(G_{\mathcal{H}}^0)_{\text{rg}} := \{v \in G^0 : s_{\mathcal{G}/\mathcal{H}}^{-1}(v) \text{ is non-empty and finite}\}.$$

Since \mathcal{H} is hereditary, if $\{v\} \in \mathcal{H}$ then we have $s_{\mathcal{G}/\mathcal{H}}^{-1}(v) = \emptyset$ and hence $v \notin (G_{\mathcal{H}}^0)_{\text{rg}}$.

LEMMA 6.8. *A hereditary ideal \mathcal{H} of \mathcal{G}^0 is saturated if and only if, whenever $v \in G_{\text{rg}}^0$ satisfies $\{v\} \notin \mathcal{H}$, we have $v \in (G_{\mathcal{H}}^0)_{\text{rg}}$.*

Proof. An element $v \in G_{\text{rg}}^0$ is in $(G_{\mathcal{H}}^0)_{\text{rg}}$ if and only if $s_{\mathcal{G}/\mathcal{H}}^{-1}(v) \subset s^{-1}(v)$ is non-empty, which occurs if and only if there is $e \in s^{-1}(v)$ with $r(e) \notin \mathcal{H}$. ■

Let \mathcal{H} be a hereditary ideal of \mathcal{G}^0 , and $U_{\mathcal{H}} \subset \Omega_{\mathcal{G}} = E(\mathcal{G})^0$ be the corresponding open subset which is hereditary by Proposition 6.6. As at the beginning of this section, we obtain a topological quiver $\mathcal{Q}(\mathcal{G})_{U_{\mathcal{H}}}$.

LEMMA 6.9. *We have $(E(\mathcal{G})_{U_{\mathcal{H}}}^0)_{\text{rg}} = (G_{\mathcal{H}}^0)_{\text{rg}}$.*

Proof. Since the image of $s_{\mathcal{Q}}|_{E(\mathcal{G})_{U_{\mathcal{H}}}^1}$ is contained in $G^0 \setminus U_{\mathcal{H}}$, we have $(E(\mathcal{G})_{U_{\mathcal{H}}}^0)_{\text{rg}} \subset G^0 \setminus U_{\mathcal{H}}$. For $v \in G^0$, $v \in U_{\mathcal{H}}$ implies $\{v\} \in \mathcal{H}$ by Lemma 6.4, and this implies $v \notin (G_{\mathcal{H}}^0)_{\text{rg}}$ as remarked after Definition 6.7. Hence we have $(G_{\mathcal{H}}^0)_{\text{rg}} \subset G^0 \setminus U_{\mathcal{H}}$. An element $v \in G^0 \setminus U_{\mathcal{H}}$ is in $(E(\mathcal{G})_{U_{\mathcal{H}}}^0)_{\text{rg}}$ if and only if $s_{\mathcal{Q}}^{-1}(v) \cap E(\mathcal{G})_{U_{\mathcal{H}}}^1$ is non-empty and compact because $\{v\}$ is open in $E(\mathcal{G})_{U_{\mathcal{H}}}^0$. Since

$$s_{\mathcal{Q}}^{-1}(v) \cap E(\mathcal{G})_{U_{\mathcal{H}}}^1 = \{(e, x) \in E(\mathcal{G})^1 : s(e) = v \text{ and } x \notin U_{\mathcal{H}}\},$$

$s_{\mathcal{Q}}^{-1}(v) \cap E(\mathcal{G})_{U_{\mathcal{H}}}^1$ is non-empty and compact if and only if

$$\{e \in \mathcal{G}^1 : s(e) = v \text{ and } \overline{r(e)} \not\subset U_{\mathcal{H}}\}$$

is non-empty and finite. This set is equal to $s_{\mathcal{G}/\mathcal{H}}^{-1}(v)$ by Lemma 6.4. Therefore an element $v \in G^0 \setminus U_{\mathcal{H}}$ is in $(E(\mathcal{G})_{U_{\mathcal{H}}}^0)_{\text{rg}}$ if and only if $v \in (G_{\mathcal{H}}^0)_{\text{rg}}$. Thus $(E(\mathcal{G})_{U_{\mathcal{H}}}^0)_{\text{rg}} = (G_{\mathcal{H}}^0)_{\text{rg}}$ as required. ■

By Lemma 6.9, the subset $(E(\mathcal{G})_{U_{\mathcal{H}}}^0)_{\text{rg}} \subset E(\mathcal{G})_{U_{\mathcal{H}}}^0$ is discrete.

DEFINITION 6.10. Let $\mathcal{G} = \{G^0, \mathcal{G}^1, r, s\}$ be an ultragraph. We say that a pair (\mathcal{H}, V) consisting of an ideal \mathcal{H} of \mathcal{G}^0 and a subset V of G^0 is *admissible* if \mathcal{H} is hereditary and saturated and $V \subset (G_{\mathcal{H}}^0)_{\text{rg}} \setminus G_{\text{rg}}^0$.

DEFINITION 6.11. For an admissible pair (\mathcal{H}, V) of an ultragraph \mathcal{G} , we define an ideal $I_{(\mathcal{H}, V)}$ of $C^*(\mathcal{G})$ to be the ideal generated by the projections

$$\{p_A : A \in \mathcal{H}\} \cup \left\{ p_v - \sum_{e \in s_{\mathcal{G}/\mathcal{H}}^{-1}(v)} s_e s_e^* : v \in V \right\}.$$

For an ideal I of $C^*(\mathcal{G})$, we define $\mathcal{H}_I := \{A \in \mathcal{G}^0 : p_A \in I\}$ and

$$V_I := \left\{ v \in (G_{\mathcal{H}_I}^0)_{\text{rg}} \setminus G_{\text{rg}}^0 : p_v - \sum_{e \in s_{\mathcal{G}/\mathcal{H}_I}^{-1}(v)} s_e s_e^* \in I \right\}.$$

THEOREM 6.12. *Let \mathcal{G} be an ultragraph. Then the correspondence $I \mapsto (\mathcal{H}_I, V_I)$ is a bijection from the set of all gauge-invariant ideals of $C^*(\mathcal{G})$ to the set of all admissible pairs of \mathcal{G} , whose inverse is given by $(\mathcal{H}, V) \mapsto I_{(\mathcal{H}, V)}$.*

Proof. By Theorem 5.2, the gauge-invariant ideals of $C^*(\mathcal{G})$ are in bijective correspondence with the gauge-invariant ideals of $C^*(\mathcal{Q}(\mathcal{G}))$. We know that the latter are indexed by admissible pairs (U, V) of $\mathcal{Q}(\mathcal{G})$ by [11, Theorem 8.22]. Proposition 6.6 and Lemma 6.9 show that $(\mathcal{H}, V) \mapsto (U_{\mathcal{H}}, V \cup (G_{\text{rg}}^0 \setminus U_{\mathcal{H}}))$ is a bijection from the set of all admissible pairs of \mathcal{G} to the one of $\mathcal{Q}(\mathcal{G})$. Thus we get bijective correspondences between

the set of all gauge-invariant ideals of $C^*(\mathcal{G})$ and the set of all admissible pairs of \mathcal{G} . By keeping track of the arguments in [11, Section 8], we see that the bijective correspondences are given by $I \mapsto (\mathcal{H}_I, V_I)$ and $(\mathcal{H}, V) \mapsto I_{(\mathcal{H}, V)}$. ■

REMARK 6.13. The theorem above naturally generalizes [1, Theorem 3.6].


7. Condition (K) In this section we define a version of condition (K) for ultragraphs, and show that this condition characterizes ultragraphs \mathcal{G} such that every ideal of $C^*(\mathcal{G})$ is gauge-invariant.

Let \mathcal{G} be an ultragraph. For $v \in G^0$, a *first-return path based at v* in \mathcal{G} is a path $\alpha = e_1 e_2 \dots e_n$ such that $s(\alpha) = v$, $v \in r(\alpha)$, and $s(e_i) \neq v$ for $i = 2, 3, \dots, n$. When α is a first-return path based at v , we say that v *hosts the first-return path α* .

Note that there is a subtlety here: a first-return path based at v may pass through other vertices $w \neq v$ more than once (that is, we may have $s(e_i) = s(e_j)$ for some $1 < i, j \leq n$ with $i \neq j$), but no edge other than e_1 may have source v .

DEFINITION 7.1. Let \mathcal{G} be an ultragraph. We say that \mathcal{G} satisfies *condition (K)* if every $v \in \mathcal{G}^0$ which hosts a first-return path hosts at least two distinct first-return paths.

EXAMPLE 7.2. The graph



satisfies condition (K) because v hosts infinitely many first-return paths $eg, efg, efgg, \dots$, and w hosts two first-return paths f and ge . Note that all first-return paths based at v except eg pass through the vertex w more than once.

PROPOSITION 7.3. Let $\mathcal{G} = (G^0, \mathcal{G}^1, r, s)$ be an ultragraph. Then every ideal of $C^*(\mathcal{G})$ is gauge-invariant if and only if \mathcal{G} satisfies condition (K).

Proof. In the same way as above, we can define first-return paths in the topological graph $E(\mathcal{G})$. It is straightforward to see that for each $v \in G^0$, first-return paths $\alpha = e_1 e_2 \dots e_n$ based at v in \mathcal{G} correspond bijectively to first-return paths

$$l = (e_1, s(e_2))(e_2, s(e_3)) \dots (e_n, s(e_1))$$

based at $v \in G^0 \subset E(\mathcal{G})^0$ in $E(\mathcal{G})$.

Recall (see [7, Definition 7.1] and the subsequent paragraph for details) that $\text{Per}(E(\mathcal{G}))$ denotes the collection of vertices $v \in E(\mathcal{G})^0$ such that v

hosts exactly one first-return path in $E(\mathcal{G})$, and v is isolated in

$$\{s_{\mathcal{Q}}(l) : l \text{ is a path in } E(\mathcal{G}) \text{ with } r_{\mathcal{Q}}(l) = v\}$$

(recall that the directions of paths are reversed when passing from the quiver $\mathcal{Q}(\mathcal{G})$ to the topological graph $E(\mathcal{G})$). We see that [7, Theorem 7.6] implies that every ideal of $\mathcal{O}(E(\mathcal{G}))$ is gauge-invariant if and only if $\text{Per}(E(\mathcal{G}))$ is empty. Since the isomorphism of $C^*(\mathcal{G})$ with $\mathcal{O}(E(\mathcal{G}))$ is gauge-equivariant, it therefore suffices to show that $\text{Per}(E(\mathcal{G}))$ is empty if and only if \mathcal{G} satisfies condition (K).

The image of $s_{\mathcal{Q}}$ is contained in the discrete set $G^0 \subset E(\mathcal{G})^0$. Thus $v \in E(\mathcal{G})^0$ belongs to $\text{Per}(E(\mathcal{G}))$ if and only if $v \in G^0 \subset E(\mathcal{G})^0$ and v hosts exactly one first-return path in $E(\mathcal{G})$. By the first paragraph of this proof, $v \in G^0 \subset E(\mathcal{G})^0$ hosts exactly one first-return path in $E(\mathcal{G})$ if and only if v hosts exactly one first-return path in \mathcal{G} . Hence $\text{Per}(E(\mathcal{G}))$ is empty if and only if \mathcal{G} satisfies condition (K). ■

References

- [1] T. Bates, J. H. Hong, I. Raeburn and W. Szymański, *The ideal structure of the C^* -algebras of infinite graphs*, Illinois J. Math. 46 (2002), 1159–1176.
- [2] J. Cuntz and W. Krieger, *A class of C^* -algebras and topological Markov chains*, Invent. Math. 56 (1980), 251–268.
- [3] R. Exel and M. Laca, *Cuntz–Krieger algebras for infinite matrices*, J. Reine Angew. Math. 512 (1999), 119–172.
- [4] C. Farthing, P. S. Muhly and T. Yeend, *Higher-rank graph C^* -algebras: an inverse semigroup and groupoid approach*, Semigroup Forum 71 (2005), 159–187.
- [5] N. J. Fowler, M. Laca and I. Raeburn, *The C^* -algebras of infinite graphs*, Proc. Amer. Math. Soc. 128 (2000), 2319–2327.
- [6] T. Katsura, *A class of C^* -algebras generalizing both graph algebras and homeomorphism C^* -algebras I, fundamental results*, Trans. Amer. Math. Soc. 356 (2004), 4287–4322.
- [7] —, *A class of C^* -algebras generalizing both graph algebras and homeomorphism C^* -algebras III, ideal structures*, Ergodic Theory Dynam. Systems 26 (2006), 1805–1854.
- [8] A. Kumjian, D. Pask and I. Raeburn, *Cuntz–Krieger algebras of directed graphs*, Pacific J. Math. 184 (1998), 161–174.
- [9] A. Kumjian, D. Pask, I. Raeburn and J. Renault, *Graphs, groupoids, and Cuntz–Krieger algebras*, J. Funct. Anal. 144 (1997), 505–541.
- [10] A. Marrero and P. S. Muhly, *Groupoid and inverse semigroup presentations of ultragraph C^* -algebras*, Semigroup Forum, to appear.
- [11] P. S. Muhly and M. Tomforde, *Topological Quivers*, Internat. J. Math. 16 (2005), 693–755.
- [12] A. L. T. Paterson, *Graph inverse semigroups, groupoids and their C^* -algebras*, J. Operator Theory 48 (2002), 645–662.
- [13] M. Tomforde, *A unified approach to Exel–Laca algebras and C^* -algebras associated to graphs*, ibid. 50 (2003), 345–368.

- [14] M. Tomforde, *Simplicity of ultragraph algebras*, Indiana Univ. Math. J. 52 (2003), 901–926.

Department of Mathematics
Keio University
Yokohama, 223-8522, Japan
E-mail: katsura@math.keio.ac.jp

Department of Mathematics
University of Iowa
Iowa City, IA 52242-1419, U.S.A.
E-mail: pmuhly@math.uiowa.edu

School of Mathematics and Applied Statistics
University of Wollongong
Wollongong, NSW 2522, Australia
E-mail: asims@uow.edu.au

Department of Mathematics
University of Houston
Houston, TX 77204-3008, U.S.A.
E-mail: tomforde@math.uh.edu

Received November 10, 2006
Revised version April 25, 2008

(6042)